

The Verlinde formula in logarithmic CFT

David Ridout^{1,2} and Simon Wood^{1,2}

¹ Department of Theoretical Physics, Research School of Physics and Engineering, Australian National University, Acton, ACT 2600, Australia

² Mathematical Sciences Institute, Australian National University, Acton, ACT 2600, Australia

E-mail: david.ridout@anu.edu.au, simon.wood@anu.edu.au

Abstract. In rational conformal field theory, the Verlinde formula computes the fusion coefficients from the modular S-transformations of the characters of the chiral algebra's representations. Generalising this formula to logarithmic models has proven rather difficult for a variety of reasons. Here, a recently proposed formalism [1] for the modular properties of certain classes of logarithmic theories is reviewed, and refined, using simple examples. A formalism addressing fusion rules in simple current extensions is also reviewed as a means to tackle logarithmic theories to which the proposed modular formalism does not directly apply.

1. Introduction

Logarithmic conformal field theory typically refers to a two-dimensional conformally invariant quantum field theory for which there exist correlation functions exhibiting logarithmic singularities. The first study of a logarithmic conformal field theory appears to be that of Rozansky and Saleur [2] where the model under consideration possesses symmetries that include the affine Kac-Moody superalgebra $\widehat{\mathfrak{gl}}(1|1)$ of central charge $c = 0$. Shortly thereafter, Gurarie noted [3] the appearance of logarithmic singularities in the $c = -2$ fermionic ghost system and showed that a sufficient condition for their presence was the non-diagonalisability of the hamiltonian $\mathbf{H} = L_0 + \bar{L}_0$. (He also seems to be responsible for the appellation “logarithmic conformal field theory”.) We refer to [4] for a recent collection of reviews on this topic.

As with rational conformal field theory, much of the analysis of logarithmic theories begins with identifying the physically relevant spectrum of representations of the chiral symmetry algebra (vertex operator algebra). Unlike the rational case, where the representations are all completely reducible, the spectrum of a logarithmic theory necessarily includes representations which are reducible, yet indecomposable.¹ In particular, the eigenvectors of \mathbf{H} always span a subrepresentation which will be proper when \mathbf{H} has non-trivial Jordan blocks. Whilst the failure of complete reducibility does not by itself guarantee Jordan blocks, hence logarithmic singularities, it seems that all the known conformal field theories whose spectrum includes reducible but indecomposable representations are logarithmic. Moreover, being logarithmic seems to be a generic property for conformal field theories, rather than a pathological one.

It is important to note, however, that logarithmic conformal field theories are necessarily non-unitary. This is not a concern for applications to statistical mechanics where the unitarity

¹ We recall that a representation is said to be indecomposable if it cannot be written as the direct sum of two non-trivial representations.

of a given lattice model is of little importance to the study of its phase transitions. Examples of lattice models whose scaling limits are known, or are believed, to be described by logarithmic conformal field theories include:

- Critical dense polymers [5], where the chiral symmetry is Virasoro (or symplectic fermions [6]) with $c = -2$.
- Critical percolation [7–10], where the symmetry is Virasoro with $c = 0$.
- Abelian sandpiles [11, 12], where the symmetry is again Virasoro with $c = -2$, but the theory appears to differ from critical dense polymers.
- The $\mathfrak{gl}(1|1)$ spin chain [7, 13, 14], where the symmetry is symplectic fermions with $c = -2$.

Non-unitarity is, of course, a concern for string-theoretic applications, but here we note that it is not the underlying conformal field theory, but only what survives the BRST quantisation (gauge-fixing) that needs to be unitary. Logarithmic theories relevant to string theory include:

- Fermionic (bc) ghosts or symplectic fermions [15, 16], where the chiral symmetry algebra is the affine Kac-Moody superalgebra $\widehat{\mathfrak{psl}}(1|1)$ with $c = -2$.
- Bosonic ($\beta\gamma$) ghosts of central charge $c = -1$ [17, 18] and $c = 2$ [19].
- Fractional level Wess-Zumino-Witten models with $\widehat{\mathfrak{sl}}(2)$ [17, 18, 20, 21] and $\widehat{\mathfrak{sl}}(3)$ [22] symmetry.
- Wess-Zumino-Witten models on Lie supergroups [2, 23–26].

There are also logarithmic conformal field theories that have been extensively studied for their own sake, in particular the $(1, p)$ triplet models [27–34] (see [35] for a proposed lattice-theoretic realisation) and their (p', p) generalisations [36–43], also known as the (W-extended) logarithmic minimal models. Recently, these ideas have been extended to $\widehat{\mathfrak{sl}}(2)$ triplets [44].

The mathematical formalism of rational conformal field theory is very rich and it is natural to try to extend the key insights to the logarithmic case. In particular, the determination of the fusion rules in rational theories is often significantly simplified through the use of the Verlinde formula [45]. This relies on computing the modular S-transformations of the characters of the irreducible representations. Unfortunately, it has been known for quite some time now [46] that the irreducible characters of a logarithmic conformal field theory need not carry a representation of the modular group, an observation which appears to completely invalidate the application of any Verlinde-type formula.

An example illustrating this dilemma is the $(1, 2)$ triplet model [27, 28] whose central charge is $c = -2$. In this model, there are four irreducible representations, $\mathbb{L}_0, \mathbb{L}_1, \mathbb{L}_{-1/8}$ and $\mathbb{L}_{3/8}$, labelled by the conformal dimensions of their highest weight states. The corresponding characters are

$$\begin{aligned} \text{ch}[\mathbb{L}_0](\tau) &= \frac{1}{2} \left(\frac{\vartheta_{1,2}(0; \tau)}{\eta(\tau)} + \eta(\tau)^2 \right), & \text{ch}[\mathbb{L}_{-1/8}](\tau) &= \frac{\vartheta_{0,2}(0; \tau)}{\eta(\tau)}, \\ \text{ch}[\mathbb{L}_1](\tau) &= \frac{1}{2} \left(\frac{\vartheta_{1,2}(0; \tau)}{\eta(\tau)} - \eta(\tau)^2 \right), & \text{ch}[\mathbb{L}_{3/8}](\tau) &= \frac{\vartheta_{2,2}(0; \tau)}{\eta(\tau)}. \end{aligned} \quad (1.1)$$

One quickly checks that it is the factors of $\eta(\tau)^2$ that lead to τ -dependent coefficients and spoil the modularity. For example,

$$\text{ch}[\mathbb{L}_0](-1/\tau) = -\frac{i\tau}{2} (\text{ch}[\mathbb{L}_0](\tau) - \text{ch}[\mathbb{L}_1](\tau)) + \frac{1}{4} (\text{ch}[\mathbb{L}_{-1/8}](\tau) - \text{ch}[\mathbb{L}_{3/8}](\tau)). \quad (1.2)$$

This can be alleviated by extending the space of characters to include a new object

$$\text{tor}(\tau) = -i\tau (\text{ch}[\mathbb{L}_0](\tau) - \text{ch}[\mathbb{L}_1](\tau)) \quad (1.3)$$

and checking that the modular S- and T-transformations preserve the space spanned by the irreducible characters and tor.

Unsurprisingly, there is a fair bit of literature addressing the issue of modularity for logarithmic conformal field theories. In particular, Miyamoto showed [47] that one is only guaranteed to have an action of the modular group on the space of torus amplitudes (one-point functions on the torus). Characters are examples of these amplitudes, as is the function tor of (1.3), so Miyamoto’s result meshes nicely with the extended character space of the (1, 2) triplet model. However, the proof requires generalising the notion of torus amplitude (and hence character) to insert arbitrary elements of the chiral symmetry algebra into the trace. While no additional elements were needed in the case of the (1, 2) triplet model, it is not clear whether this will be true more generally. If not, then evaluating these refined characters and torus amplitudes leads to severe computational difficulties in all but the simplest examples. It also leaves one wondering how such a result helps in determining modular invariant bulk partition functions (which are, after all, sesquilinear combinations of characters).

Finding modular invariants therefore seems to require new insight [48], but one can instead ask a different question. Given a logarithmic conformal field theory, is there a variant of the Verlinde formula which can be used to compute the fusion rules efficiently? The technical answer to this question is “no”, as long as we assume that the input to the Verlinde formula is the S-transformation of characters, because characters cannot distinguish between a representation and the direct sum of its irreducible constituents (composition factors). Instead, the best that we can hope for is a Verlinde-type formula that determines the *Grothendieck* fusion rules that are obtained from the genuine fusion rules by identifying representations with the same character.²

Is there a Verlinde formula that computes the Grothendieck fusion rules? For the (1, p) triplet models, the answer is “yes” [49]. Here, one computes the modular S-matrix S on the basis of irreducible characters directly, so that the entries are τ -dependent in general. But then, the action is twisted by a carefully chosen τ -dependent automorphy factor A such that the product $S' = AS$ is both independent of τ and, with T , defines a representation of the modular group on character space. Remarkably, this matrix S' successfully block-diagonalises the Grothendieck fusion rules in a Verlinde-type formula given in [49, p717]. Most importantly, the entries of the block-diagonal matrix are also expressed in terms of the entries of S' .

Admittedly, this formula is rather more complicated than the Verlinde formula of rational conformal field theory. However, this is to be expected from the non-diagonalisability of the fusion matrices in the logarithmic case. But, there seems to be little understanding of why twisting by this automorphy factor A is required, whether there are other possibilities for A , and whether, for more general models, the existence of such a factor is always guaranteed. As far as we are aware, this type of Verlinde formula has not been generalised beyond the (1, p) triplet models. Moreover, there are now several other proposals [30, 50–52] addressing seemingly different Verlinde formulae for these theories.

Rather than try to compare and contrast these proposals, which would be a worthy goal in itself, we shall instead turn to a rather different formalism which has recently been introduced [1] for modular properties and Verlinde formulae in a more general class of logarithmic conformal field theories. One difference between this proposal and those mentioned above is that it does not directly apply to the triplet models. However, this is not a serious restriction because the triplet models have close relatives to which the new formalism does apply. We shall explain how this allows us to compute Grothendieck fusion rules for the triplet models in Section 3.

² There is a subtle mathematical question that we are overlooking here of whether the notion of Grothendieck fusion is well-defined. See [38] for an example where it is not.

2. Modularity for logarithmic CFT

The formalism discussed in this section arises from the observation [1, 25] that the majority of the logarithmic conformal field theories that have been adequately analysed to date all share certain key features. Moreover, the known exceptions may all be understood as simple current extensions of theories with the desired features. We will therefore describe these features before explaining, with the aid of examples, how they allow one to straightforwardly compute modular transformations and apply the Verlinde formula to obtain Grothendieck fusion coefficients.

The formalism centres [1] about the identification of a collection of representations of the chiral algebra (vertex operator algebra) that we call the *standard modules*. These standard modules and their characters should have the following properties:

- (1) The characters of the standard modules are parametrised by a measurable space (M, μ) .
- (2) The irreducible standard modules are called *typical*. The complement of the subset of M corresponding to the typical characters has measure zero with respect to μ .
- (3) The only indecomposable module in the spectrum having a typical module (irreducible standard) as a composition factor (irreducible subquotient) is that typical module.³
- (4) The characters of the standard modules form a (topological) basis for the space of all characters. In particular, the standard module characters are linearly independent.
- (5) The characters of the standard modules span a representation of the modular group $\mathrm{SL}(2; \mathbb{Z})$. In the standard basis, the S-transformation defines a symmetric, unitary operator whose square may be identified with conjugation.
- (6) Combining the S-transformation with the standard Verlinde formula leads to the Grothendieck fusion rules (with non-negative integer coefficients).

Any module in the spectrum that is not a direct sum of typical modules is called *atypical*, the nomenclature here being inspired by that of Kac for Lie superalgebra representations. We remark that there will usually be more than one inequivalent standard module for each atypical parameter $m \in M$, but each of these atypical standard modules will have the same character.

We will not give any general means for identifying the standard modules in a given theory beyond the obvious need to satisfy the above requirements; in examples, this identification is often clear. In this vein, we mention that verifying (3) rigorously is almost always too difficult at present, so we usually have to content ourselves with conjecturing that it holds. We also remark that (4) is generally seen to be satisfied by the existence of resolutions for the irreducible atypical modules in terms of atypical standard modules. The Euler-Poincaré principle then expresses the character of each irreducible as an (infinite) alternating sum of standard characters.

The main observation that we want to emphasise is that the typical modules behave, for all intents and purposes, as if they were in a rational conformal field theory. Indeed, rational theories fit into the above formalism as the special case in which all the standard modules are typical. The logarithmic nature of the theory is therefore concentrated in the atypical modules and these are required to form a set of measure zero in character-space. This means that, heuristically at least, any potential problems that the logarithmic structure would create are measure-theoretically void. While there is an open question as to how one identifies the physically relevant measure in general, we will see that in examples, the natural choice works.

Speaking of which, the simplest (non-rational) example illustrating this formalism is the (uncompactified) free boson. However, as this model is not logarithmic, we will refer the reader to [1, Sec. 1.2] and instead turn to a closely related logarithmic conformal field theory.

³ Another way of stating this would be to say that typical modules are both projective and injective in the physically relevant category of modules. However, this is extremely difficult to check in practice, partly because it is not clear how one actually identifies the physically relevant category.

Example 1 (The singlet model [1, 53]). The fermionic bc ghost system at $c = -2$ contains the symplectic fermions algebra as the subalgebra generated by b and ∂c . This subalgebra in turn contains a subalgebra corresponding to the states of zero ghost number. This is the $(1, 2)$ singlet algebra and it is generated by the energy-momentum tensor and a single Virasoro primary W of conformal dimension 3.

- (1) The singlet algebra possesses a continuous family of highest weight modules \mathcal{F}_λ , with $\lambda \in \mathbb{R}$. These are the standard modules. Their characters count the eigenvalues of L_0 , but these do not distinguish conjugate modules. We therefore augment them by counting the eigenvalues of a shifted ghost number, the zero mode of $G = :bc: - \frac{1}{2}$, and the central element $\mathbf{1}$:

$$\text{ch}[\mathcal{F}_\lambda](y; z; q) = \text{tr}_{\mathcal{F}_\lambda} y^{\mathbf{1}} z^{G_0} q^{L_0+1/12} = \frac{yz^{\lambda-1/2} q^{(\lambda-1/2)^2/2}}{\eta(q)}. \quad (2.1)$$

- (2) The standard modules are irreducible if and only if $\lambda \notin \mathbb{Z}$. We therefore equip \mathbb{R} with Lebesgue measure μ , noting that $\mu(\mathbb{Z}) = 0$.
- (3) We cannot prove that the typicals are projective and injective, but it seems to be true.
- (4) The standard characters (2.1) are clearly linearly independent. The atypical standard modules \mathcal{F}_λ , with $\lambda \in \mathbb{Z}$, are built from two composition factors (irreducible subquotients). The structure is given by the following short exact sequences:⁴

$$0 \longrightarrow \mathcal{L}_\lambda \longrightarrow \mathcal{F}_\lambda \longrightarrow \mathcal{L}_{\lambda-1} \longrightarrow 0. \quad (2.2)$$

We then splice (see [1, App. A.2] for example) these exact sequences to obtain resolutions:

$$\cdots \longrightarrow \mathcal{F}_{\lambda+3} \longrightarrow \mathcal{F}_{\lambda+2} \longrightarrow \mathcal{F}_{\lambda+1} \longrightarrow \mathcal{L}_\lambda \longrightarrow 0. \quad (2.3)$$

This is also an exact sequence, so the Euler-Poincaré principle leads to the character formula

$$\text{ch}[\mathcal{L}_\lambda] = \sum_{n=1}^{\infty} (-1)^{n-1} \text{ch}[\mathcal{F}_{\lambda+n}] \quad (\lambda \in \mathbb{Z}), \quad (2.4)$$

verifying that the standard characters form a topological basis of the space of all characters.

- (5) The standard singlet characters span a representation of the modular group whose dimension is uncountably-infinite. The S-transformation is essentially a Fourier transform:

$$\begin{aligned} \text{ch}[\mathcal{F}_\lambda]\left(\theta - \frac{\zeta^2}{2\tau}; \frac{\zeta}{\tau}; -\frac{1}{\tau}\right) &= \int_{\mathbb{R}} \mathbf{S}[\mathcal{F}_\lambda \rightarrow \mathcal{F}_{\lambda'}] \text{ch}[\mathcal{F}_{\lambda'}](\theta; \zeta; \tau) d\lambda', \\ \mathbf{S}[\mathcal{F}_\lambda \rightarrow \mathcal{F}_{\lambda'}] &= e^{-2\pi i(\lambda-1/2)(\lambda'-1/2)}. \end{aligned} \quad (2.5)$$

This is symmetric and unitary and it squares to the conjugation permutation $\lambda \leftrightarrow 1 - \lambda$:

$$\mathbf{S}[\mathcal{F}_\lambda \rightarrow \mathcal{F}_{\lambda'}] = \mathbf{S}[\mathcal{F}_{\lambda'} \rightarrow \mathcal{F}_\lambda], \quad (2.6a)$$

$$\int_{\mathbb{R}} \mathbf{S}[\mathcal{F}_\lambda \rightarrow \mathcal{F}_{\lambda'}] \mathbf{S}[\mathcal{F}_{\lambda''} \rightarrow \mathcal{F}_{\lambda'}]^* d\lambda' = \delta(\lambda - \lambda''), \quad (2.6b)$$

$$\int_{\mathbb{R}} \mathbf{S}[\mathcal{F}_\lambda \rightarrow \mathcal{F}_{\lambda'}] \mathbf{S}[\mathcal{F}_{\lambda'} \rightarrow \mathcal{F}_{\lambda''}] d\lambda' = \delta(\lambda + \lambda'' - 1). \quad (2.6c)$$

The atypical S-transformations are now obtained from the character formula (2.4):

$$\mathbf{S}[\mathcal{L}_\lambda \rightarrow \mathcal{F}_{\lambda'}] = \sum_{n=1}^{\infty} (-1)^{n-1} \mathbf{S}[\mathcal{F}_{\lambda+n} \rightarrow \mathcal{F}_{\lambda'}] = \frac{e^{-2\pi i\lambda(\lambda'-1/2)}}{2 \cos(\pi(\lambda' - 1/2))}. \quad (2.7)$$

We note that these elements *diverge* precisely at the atypical parameters $\lambda' \in \mathbb{Z}$.

⁴ Exactness means that at each position $\cdots \xrightarrow{f} M \xrightarrow{g} \cdots$ in the sequence, the image of f coincides with the kernel of g . If $0 \rightarrow L \rightarrow M \rightarrow N \rightarrow 0$ is exact, then L is a submodule of M and M/L is isomorphic to N .

(6) The (Grothendieck) fusion rules are given by the following version of the Verlinde formula:

$$\begin{aligned} \begin{bmatrix} \mathcal{F}_{\lambda''} \\ \mathcal{F}_{\lambda} & \mathcal{F}_{\lambda'} \end{bmatrix} &= \int_{\mathbb{R}} \frac{\mathcal{S}[\mathcal{F}_{\lambda} \rightarrow \mathcal{F}_{\lambda''}] \mathcal{S}[\mathcal{F}_{\lambda'} \rightarrow \mathcal{F}_{\lambda''}] \mathcal{S}[\mathcal{F}_{\lambda''} \rightarrow \mathcal{F}_{\lambda''}]^*}{\mathcal{S}[\mathcal{L}_0 \rightarrow \mathcal{F}_{\lambda''}]} d\lambda'', \\ \text{ch}[\mathcal{F}_{\lambda}] \times \text{ch}[\mathcal{F}_{\lambda'}] &= \int_{\mathbb{R}} \begin{bmatrix} \mathcal{F}_{\lambda''} \\ \mathcal{F}_{\lambda} & \mathcal{F}_{\lambda'} \end{bmatrix} \text{ch}[\mathcal{F}_{\lambda''}] d\lambda''. \end{aligned} \quad (2.8)$$

Note that \mathcal{L}_0 is the vacuum module. There are atypical analogues obtained by replacing \mathcal{F}_{λ} or $\mathcal{F}_{\lambda'}$ (but never $\mathcal{F}_{\lambda''}$) by atypical modules. We thereby obtain the Grothendieck fusion rules

$$\begin{aligned} \text{ch}[\mathcal{L}_{\lambda}] \times \text{ch}[\mathcal{L}_{\lambda'}] &= \text{ch}[\mathcal{L}_{\lambda+\lambda'}], & \text{ch}[\mathcal{L}_{\lambda}] \times \text{ch}[\mathcal{F}_{\lambda'}] &= \text{ch}[\mathcal{F}_{\lambda+\lambda'}], \\ \text{ch}[\mathcal{F}_{\lambda}] \times \text{ch}[\mathcal{F}_{\lambda'}] &= \text{ch}[\mathcal{F}_{\lambda+\lambda'}] + \text{ch}[\mathcal{F}_{\lambda+\lambda'-1}], \end{aligned} \quad (2.9)$$

confirming that the Grothendieck fusion coefficients are all non-negative integers. The atypical-atypical rule lifts to a genuine fusion rule because the result is an irreducible character that determines the module uniquely. The same is true for the atypical-typical rule because $\lambda + \lambda' \notin \mathbb{Z}$. The typical-typical rule, however, can only be lifted to a direct sum when $\lambda + \lambda'$ is typical. When $\lambda + \lambda' \in \mathbb{Z}$, we expect that the fusion product is a staggered module in the sense of [1, 54].⁵

The observed divergence (2.7) of the atypical S-transformation kernel at atypical parameters seems to be a general feature of logarithmic theories. For this reason, it is absolutely vital that the atypical standards correspond to a set of measure zero in parameter space. We will develop a method to deal with the case when the atypical parameters have positive measure in Section 3. Before turning to this, however, we detail another, slightly more involved, logarithmic example.

Example 2 (Bosonic ghosts [19]). The $\beta\gamma$ ghost system with $c = 2$ possesses a unique highest weight representation, the vacuum module \mathcal{V} . This module is irreducible. However, the ghost algebra admits a family of spectral flow automorphisms σ^{ℓ} which yields a family of mutually inequivalent representations $\sigma^{\ell}(\mathcal{V})$, $\ell \in \mathbb{Z}$. These are *not* the standard modules of the theory because their characters do *not* span a representation of the modular group.

(1) The ghost algebra also admits a continuous family of *parabolic* Verma modules \mathcal{W}_j , with $0 < j < 1$. Moreover, there exist two additional inequivalent indecomposable modules, \mathcal{W}_0^+ and \mathcal{W}_0^- , characterised by the short exact sequences

$$0 \longrightarrow \mathcal{V} \longrightarrow \mathcal{W}_0^+ \longrightarrow \sigma^{-1}(\mathcal{V}) \longrightarrow 0, \quad 0 \longrightarrow \sigma^{-1}(\mathcal{V}) \longrightarrow \mathcal{W}_0^- \longrightarrow \mathcal{V} \longrightarrow 0. \quad (2.10)$$

The spectral flow images $\sigma^{\ell}(\mathcal{W}_j)$, together with the $\sigma^{\ell}(\mathcal{W}_0^{\pm})$, are the standard modules. The standard characters again involve the ghost number, obtained from the zero mode of $J = : \beta\gamma : ,$ as well as the central element $\mathbf{1}$:

$$\text{ch}[\sigma^{\ell}(\mathcal{W}_j)](y; z; q) = \text{tr}_{\sigma^{\ell}(\mathcal{W}_j)} y^{\mathbf{1}} z^{J_0} q^{L_0 - 1/12} = \frac{y z^j q^{\ell j + \ell(\ell-1)/2}}{\eta(q)^2} \sum_{n \in \mathbb{Z}} z^n q^{n\ell}. \quad (2.11)$$

The common character of $\sigma^{\ell}(\mathcal{W}_0^+)$ and $\sigma^{\ell}(\mathcal{W}_0^-)$ is also given by this formula (with j set to 0). The standard characters are thus naturally parametrised by $j \in \mathbb{R}/\mathbb{Z} \cong \mathbb{S}^1$ and $\ell \in \mathbb{Z}$. We equip this space with the product μ of the standard Haar measure on \mathbb{S}^1 and counting measure on \mathbb{Z} .

⁵ Unfortunately, nobody seems to have computed fusion products for the (1, 2) singlet algebra. However, this expectation is consistent with the known fusion rules for the (1, 2) *triplet* algebra [27] as we shall see.

- (2) The $\sigma^\ell(\mathcal{W}_j)$, with $j \neq 0$, are irreducible. These are the typical modules: $\mu(\{0\} \times \mathbb{Z}) = 0$.
(3) Again, we content ourselves with conjecturing that the typicals are projective and injective.
(4) Splicing (2.10) to obtain resolutions, we arrive at the atypical character formula

$$\text{ch}[\sigma^\ell(\mathcal{V})] = \sum_{n=1}^{\infty} (-1)^{n-1} \text{ch}[\sigma^{\ell+n}(\mathcal{W}_0^+)], \quad (2.12)$$

which confirms that the standard characters form a topological basis.

- (5) The modular transformations take a rather more complicated form for this model:

$$\begin{aligned} \mathbf{S}: (\theta; \zeta; \tau) &\longrightarrow \left(\theta + \frac{\zeta^2}{2\tau} - \frac{\zeta}{2\tau} + \frac{\zeta}{2} + \frac{1}{2\pi} \left(\arg \tau - \frac{\pi}{2} \right); \frac{\zeta}{\tau}; -\frac{1}{\tau} \right), \\ \mathbf{T}: (\theta; \zeta; \tau) &\longrightarrow \left(\theta + \frac{1}{12}; \zeta; \tau + 1 \right). \end{aligned} \quad (2.13)$$

Nevertheless, they determine a representation of $\text{SL}(2; \mathbb{Z})$ on the span of the standard characters with S-transformation

$$\begin{aligned} \text{ch}[\sigma^\ell(\mathcal{W}_j)](\mathbf{S}(\theta; \zeta; \tau)) &= \sum_{\ell' \in \mathbb{Z}} \int_{\mathbb{R}/\mathbb{Z}} \mathbf{S}[\sigma^\ell(\mathcal{W}_j) \rightarrow \sigma^{\ell'}(\mathcal{W}_{j'})] \text{ch}[\sigma^{\ell'}(\mathcal{W}_{j'})](\theta; \zeta; \tau) dj', \\ \mathbf{S}[\sigma^\ell(\mathcal{W}_j) \rightarrow \sigma^{\ell'}(\mathcal{W}_{j'})] &= (-1)^{\ell+\ell'} e^{-2\pi i(\ell j' + \ell' j)}. \end{aligned} \quad (2.14)$$

It is easy to check that \mathbf{S} is symmetric, unitary and squares to conjugation. The atypical S-transformation kernel

$$\mathbf{S}[\sigma^\ell(\mathcal{V}) \rightarrow \sigma^{\ell'}(\mathcal{W}_{j'})] = (-1)^{\ell+\ell'+1} \frac{e^{-2\pi i(\ell+1/2)j'}}{e^{i\pi j'} - e^{-i\pi j'}}, \quad (2.15)$$

again diverges at the atypical point $j' = 0$.

- (6) The obvious Verlinde formula (with sums over $\ell \in \mathbb{Z}$ and integrals over $j \in \mathbb{R}/\mathbb{Z}$) gives the following Grothendieck fusion rules:

$$\text{ch}[\sigma^\ell(\mathcal{V})] \times \text{ch}[\sigma^{\ell'}(\mathcal{V})] = \text{ch}[\sigma^{\ell+\ell'}(\mathcal{V})], \quad (2.16a)$$

$$\text{ch}[\sigma^\ell(\mathcal{V})] \times \text{ch}[\sigma^{\ell'}(\mathcal{W}_{j'})] = \text{ch}[\sigma^{\ell+\ell'}(\mathcal{W}_{j'})], \quad (2.16b)$$

$$\text{ch}[\sigma^\ell(\mathcal{W}_j)] \times \text{ch}[\sigma^{\ell'}(\mathcal{W}_{j'})] = \text{ch}[\sigma^{\ell+\ell'}(\mathcal{W}_{j+j'})] + \text{ch}[\sigma^{\ell+\ell'-1}(\mathcal{W}_{j+j'})]. \quad (2.16c)$$

The coefficients are clearly non-negative in each case.

We close this section by noting that this formalism has been extensively tested on other logarithmic conformal field theories including the fractional level $\widehat{\mathfrak{sl}}(2)$ models [55, 56], the $\widehat{\mathfrak{gl}}(1|1)$ model [25, 57] and its Takiff extension [58], the logarithmic minimal models [59] and their $N = 1$ extensions [60], and the (p', p) singlet theories [53]. The Takiff model is of interest because it demonstrates that the formalism can naturally accommodate different degrees of atypicality. The singlet theories with $p, p' \geq 2$ are interesting because it is known [38] that their fusion product does not induce a well-defined product on the Grothendieck group because of the poor behaviour (non-rigidity) of the minimal model representations. The formalism deals with this naturally, and rather effectively, by regarding the minimal model characters as being zero.⁶

⁶ One is therefore led to wonder whether the minimal model representations should be excluded from the physically relevant category. They do not seem to appear as direct summands in the bulk, boundary or defect sectors. However, excluding them would amount to taking the physical category to be non-abelian.

3. Fusion Rules for Simple Current Extensions

We recall that a simple current may be generally defined as an indecomposable representation for which fusing with the simple current defines a permutation on the set of all indecomposables. For the purposes of this article, we shall simplify this definition by replacing “indecomposable” by “irreducible”, as in [61, 62]. Examples of such simple currents then include the \mathcal{L}_λ , with $\lambda \in \mathbb{Z}$, for the $(1, 2)$ singlet model (Example 1) and the $\sigma^\ell(\mathcal{V})$, with $\ell \in \mathbb{Z}$, for the $c = 2$ bosonic ghost system (Example 2). In both cases, the simple currents are atypical irreducibles.

If the vacuum module is also irreducible, as is the case for both examples considered in Section 2, then for every simple current \mathcal{J} , there exists a simple current \mathcal{J}^{-1} such that their fusion gives the vacuum module, that is, \mathcal{J} has a fusion inverse. Any given simple current (more generally, any set of simple currents) then generates a group \mathbf{G} of simple currents under the fusion operation. This group then generates an extended symmetry algebra in the sense that the representation $\bigoplus_{\mathcal{J} \in \mathbf{G}} \mathcal{J}$ carries the structure of a vertex operator algebra.⁷ Moreover, every representation \mathcal{M} of the original symmetry algebra determines⁸ a representation of the extended symmetry algebra on $\bigoplus_{\mathcal{J} \in \mathbf{G}} (\mathcal{J} \times \mathcal{M})$.

Mathematically, this may be formalised using the notions of restriction and induction [66, 67]: Let \mathbf{A} denote the original vertex operator algebra, \mathbf{B} its simple current extension by \mathbf{G} , and $\otimes_{\mathbf{A}}$ and $\otimes_{\mathbf{B}}$ their fusion products. Then, restricting \mathbf{B} (interpreted as its vacuum module) to an \mathbf{A} -module and inducing \mathbf{A} (also interpreted as its vacuum module) to a \mathbf{B} -module amounts to

$$\mathbf{B} \downarrow \cong \bigoplus_{\mathcal{J} \in \mathbf{G}} \mathcal{J}, \quad \mathbf{A} \uparrow = \mathbf{B} \otimes_{\mathbf{A}} \mathbf{A} \cong \mathbf{B}. \quad (3.1)$$

Moreover, we may induce an \mathbf{A} -module \mathcal{M} to a \mathbf{B} -module $\mathcal{M} \uparrow$, obtaining

$$\mathcal{M} \uparrow = \mathbf{B} \otimes_{\mathbf{A}} \mathcal{M} \quad \Rightarrow \quad (\mathcal{M} \uparrow) \downarrow \cong \bigoplus_{\mathcal{J} \in \mathbf{G}} (\mathcal{J} \otimes_{\mathbf{A}} \mathcal{M}), \quad (3.2)$$

as before. This formalisation has a practical benefit as well. The associativity of fusion products allows one to compute the fusion rules for the simple current extension \mathbf{B} from those of \mathbf{A} :

$$\begin{aligned} \mathcal{M} \uparrow \otimes_{\mathbf{B}} \mathcal{N} \uparrow &= (\mathbf{B} \otimes_{\mathbf{A}} \mathcal{M}) \otimes_{\mathbf{B}} (\mathbf{B} \otimes_{\mathbf{A}} \mathcal{N}) \cong ((\mathbf{B} \otimes_{\mathbf{A}} \mathcal{M}) \otimes_{\mathbf{B}} \mathbf{B}) \otimes_{\mathbf{A}} \mathcal{N} \\ &\cong (\mathbf{B} \otimes_{\mathbf{A}} \mathcal{M}) \otimes_{\mathbf{A}} \mathcal{N} \cong \mathbf{B} \otimes_{\mathbf{A}} (\mathcal{M} \otimes_{\mathbf{A}} \mathcal{N}) = (\mathcal{M} \otimes_{\mathbf{A}} \mathcal{N}) \uparrow. \end{aligned} \quad (3.3)$$

Here, we remark that inducing an irreducible \mathbf{A} -module \mathcal{M} results in an irreducible \mathbf{B} -module $\mathcal{M} \uparrow$. This is a special property of (invertible) simple current extensions that follows from (3.2) because each direct summand $\mathcal{J} \otimes_{\mathbf{A}} \mathcal{M}$ is an irreducible that is obtained from any other direct summand by acting with one of the simple currents in \mathbf{G} , thus with an element of \mathbf{B} .

Of course, one needs to justify why the fusion product of a vertex operator algebra may be regarded as a tensor product that is permeable to the algebra action. That fusion constitutes an abstract tensor product is, at least for C_2 -cofinite vertex operator algebras, one of the great results of Lepowsky’s group (see [68] for a review). While the logarithmic theories we study here are not C_2 -cofinite in general, it is very reasonable to assume that fusion still has the properties of a tensor product on the physical category of representations. The permeability, on the other hand, reflects the locality requirement for fields corresponding to the algebra and its modules. More precisely, if $a(z)$ is a field of \mathbf{A} and $\phi(z)$ and $\psi(z)$ are fields corresponding to elements of the \mathbf{A} -modules \mathcal{M} and \mathcal{N} , respectively, then locality and associativity give

$$\begin{aligned} (a(z) \phi(x)) \psi(w) &= (\phi(x) a(z)) \psi(w) = \phi(x) (a(z) \psi(w)) \\ \iff a \cdot \phi \otimes_{\mathbf{A}} \psi &= \phi \cdot a \otimes_{\mathbf{A}} \psi = \phi \otimes_{\mathbf{A}} a \cdot \psi. \end{aligned} \quad (3.4)$$

⁷ There are some subtleties to this statement, however; see [21, 64, 65] for example.

⁸ For simplicity, and because our examples do not require it, we ignore the issue of *fixed points* here, see [63].

We conclude by computing the Grothendieck fusion rules of the $(1, 2)$ triplet model as an application of this simple current formalism. Recall from Equation (1.2) that the irreducible characters do not span a representation of the modular group. From the point of view of the modular formalism of Section 2, one identifies candidates for the standard modules: The typical irreducibles $\mathbb{L}_{-1/8}$ and $\mathbb{L}_{3/8}$ and the atypical indecomposables \mathbb{A}_0 and \mathbb{A}_1 are characterised by

$$0 \longrightarrow \mathbb{L}_0 \longrightarrow \mathbb{A}_0 \longrightarrow \mathbb{L}_1 \longrightarrow 0, \quad 0 \longrightarrow \mathbb{L}_1 \longrightarrow \mathbb{A}_1 \longrightarrow \mathbb{L}_0 \longrightarrow 0. \quad (3.5)$$

Unfortunately, the standard characters do not span a representation of the modular group either. We remark that as $\text{ch}[\mathbb{A}_0] = \text{ch}[\mathbb{A}_1]$, the atypical characters have measure $\frac{1}{3}$ with respect to the natural measure (counting measure) on the parametrisation space $M = \{-\frac{1}{8}, 0, \frac{3}{8}\}$.

As mentioned above, (Grothendieck) fusion rules for the $(1, 2)$ triplet model are nevertheless easily derived by regarding this theory as the simple current extension of the $(1, 2)$ singlet model. The group \mathbb{G} of simple currents is generated by the irreducible singlet representation $\mathcal{J} = \mathcal{L}_2$ (or \mathcal{L}_{-2}) [1] whose highest weight state has conformal dimension 3. We can therefore apply the above simple current formalism with \mathbb{A} and \mathbb{B} being the singlet and triplet vertex operator algebras, respectively. It is straightforward to identify the triplet irreducibles as induced singlet modules (see Example 1):

$$\begin{aligned} \mathbb{L}_0 &= \mathcal{L}_0 \uparrow, & \mathbb{L}_1 &= \mathcal{L}_1 \uparrow, & \mathbb{L}_{-1/8} &= \mathcal{F}_{1/2} \uparrow, & \mathbb{L}_{3/8} &= \mathcal{F}_{-1/2} \uparrow, \\ \mathbb{L}_0 \downarrow &= \bigoplus_{n \in \mathbb{Z}} \mathcal{L}_{2n}, & \mathbb{L}_1 \downarrow &= \bigoplus_{n \in \mathbb{Z}} \mathcal{L}_{2n+1}, & \mathbb{L}_{-1/8} \downarrow &= \bigoplus_{n \in \mathbb{Z}} \mathcal{F}_{2n+1/2}, & \mathbb{L}_{3/8} \downarrow &= \bigoplus_{n \in \mathbb{Z}} \mathcal{F}_{2n-1/2}. \end{aligned} \quad (3.6)$$

The remaining singlet typicals \mathcal{F}_λ can likewise be induced, but they turn out to define *twisted* representations on which the triplet fields act with non-trivial monodromy.

The singlet fusion rules derived in Example 1 may now be lifted to the triplet using (3.3). For example,

$$\mathbb{L}_1 \otimes_{\mathbb{B}} \mathbb{L}_1 = \mathcal{L}_1 \uparrow \otimes_{\mathbb{B}} \mathcal{L}_1 \uparrow = (\mathcal{L}_1 \otimes_{\mathbb{A}} \mathcal{L}_1) \uparrow = \mathcal{L}_2 \uparrow = \mathbb{L}_0. \quad (3.7a)$$

In the same manner, we compute

$$\mathbb{L}_1 \otimes_{\mathbb{B}} \mathbb{L}_{-1/8} = \mathbb{L}_{3/8}, \quad \mathbb{L}_1 \otimes_{\mathbb{B}} \mathbb{L}_{3/8} = \mathbb{L}_{-1/8}. \quad (3.7b)$$

These results agree of course with direct, though significantly more difficult, computations [27]. For the fusion rules of $\mathbb{L}_{-1/8}$ and $\mathbb{L}_{3/8}$, the corresponding singlet rules were only derived at the Grothendieck level, so we may only conclude that

$$\text{ch}[\mathbb{L}_h] \otimes_{\mathbb{B}} \text{ch}[\mathbb{L}_{h'}] = 2 \text{ch}[\mathbb{L}_0] + 2 \text{ch}[\mathbb{L}_1], \quad (h, h' \in \{-\frac{1}{8}, \frac{3}{8}\}). \quad (3.8)$$

This result is also consistent with the known fusion rules [27] where the fusion products are known to be indecomposable staggered modules formed by glueing two copies of \mathbb{L}_0 and two copies of \mathbb{L}_1 together.

This example illustrates how one can combine the standard module formalism of Section 2 with the simple current formalism of Section 3 to rather effortlessly deduce the (Grothendieck) fusion rules of logarithmic conformal field theories. The results have been generalised to all the (p', p) triplet models [53] and are in complete agreement with the fusion computations that have been computed, as well as those that have only been conjectured, for these theories. We conclude by remarking that we are unaware of theories to which this combined formalism cannot be applied. It would be very interesting to study an example of such a theory.

Acknowledgements

We thank Jürgen Fuchs and Aliosha Semikhatov for discussions on the Verlinde formula for the $(1, p)$ triplet models, Matthias Gaberdiel for helpful correspondence concerning the modular properties of the triplet characters, and Antun Milas for pointing out a mistake in an earlier version. DR's research is supported by the Australian Research Council Discovery Project DP1093910. SW's work is supported by the Australian Research Council Discovery Early Career Researcher Award DE140101825.

References

- [1] Creutzig T and Ridout D 2013 *J. Phys.* **A46** 494006 arXiv:1303.0847 [hep-th]
- [2] Rozansky L and Saleur H 1992 *Nucl. Phys.* **B376** 461–509
- [3] Gurarie V 1993 *Nucl. Phys.* **B410** 535–549 arXiv:hep-th/9303160
- [4] Gainutdinov A, Ridout D and Runkel I (Editors) 2013 *J. Phys.* **A46** 490301
- [5] Pearce P and Rasmussen J 2007 *J. Stat. Mech.* **0702** P02015 arXiv:hep-th/0610273
- [6] Morin-Duchesne A, Pearce P and Rasmussen J 2013 *Nucl. Phys.* **B874** 312–357 arXiv:1303.4895 [hep-th]
- [7] Read N and Saleur H 2007 *Nucl. Phys.* **B777** 316–351 arXiv:hep-th/0701117
- [8] Rasmussen J and Pearce P 2007 *J. Stat. Mech.* **0709** P09002 arXiv:0706.2716 [hep-th]
- [9] Mathieu P and Ridout D 2007 *Phys. Lett.* **B657** 120–129 arXiv:0708.0802 [hep-th]
- [10] Ridout D 2009 *Nucl. Phys.* **B810** 503–526 arXiv:0808.3530 [hep-th]
- [11] Jeng M, Piroux G and Ruelle P 2006 *J. Stat. Mech.* **0610** P10015 arXiv:cond-mat/0609284
- [12] Ruelle P 2013 *J. Phys.* **A46** 494014 arXiv:1303.4310 [hep-th]
- [13] Gainutdinov A, Read N and Saleur H 2013 *Nucl. Phys.* **B871** 245–288 arXiv:1112.3403 [hep-th]
- [14] Gainutdinov A, Jacobsen J, Read N, Saleur H and Vasseur R 2013 *J. Phys.* **A46** 494012 arXiv:1303.2082 [hep-th]
- [15] Kausch H Curiosities at $c = -2$ arXiv:hep-th/9510149
- [16] Kausch H 2000 *Nucl. Phys.* **B583** 513–541 arXiv:hep-th/0003029
- [17] Lesage F, Mathieu P, Rasmussen J and Saleur H 2004 *Nucl. Phys.* **B686** 313–346 arXiv:hep-th/0311039
- [18] Ridout D 2011 *Nucl. Phys.* **B848** 216–250 arXiv:1012.2905 [hep-th]
- [19] Ridout D and Wood S Bosonic Ghosts at $c = 2$ as a Logarithmic CFT arXiv:1408.4185 [hep-th]
- [20] Gaberdiel M 2001 *Nucl. Phys.* **B618** 407–436 arXiv:hep-th/0105046
- [21] Ridout D 2009 *Nucl. Phys.* **B814** 485–521 arXiv:0810.3532 [hep-th]
- [22] Adamović D A Realization of Certain Modules for the $N = 4$ Superconformal Algebra and the Affine Lie Algebra $A_2^{(1)}$ arXiv:1407.1527 [math.QA]
- [23] Saleur H and Schomerus V 2006 *Nucl. Phys.* **B734** 221–245 arXiv:hep-th/0510032
- [24] Gotz G, Quella T and Schomerus V 2007 *JHEP* **0703** 003 arXiv:hep-th/0610070
- [25] Creutzig T and Ridout D 2013 *Nucl. Phys.* **B872** 348–391 arXiv:1107.2135 [hep-th]
- [26] Quella T and Schomerus V 2013 *J. Phys.* **A46** 494010 arXiv:1307.7724 [hep-th]
- [27] Gaberdiel M and Kausch H 1996 *Phys. Lett.* **B386** 131–137 arXiv:hep-th/9606050
- [28] Gaberdiel M and Kausch H 1999 *Nucl. Phys.* **B538** 631–658 arXiv:hep-th/9807091
- [29] Feigin B, Gainutdinov A, Semikhatov A and Tipunin I Y 2006 *Comm. Math. Phys.* **265** 47–93 arXiv:hep-th/0504093
- [30] Gaberdiel M and Runkel I 2008 *J. Phys.* **A41** 075402 arXiv:0707.0388 [hep-th]
- [31] Adamović D and Milas A 2008 *Adv. Math.* **217** 2664–2699 arXiv:0707.1857 [math.QA]
- [32] Pearce P, Rasmussen J and Ruelle P 2008 *J. Phys.* **A41** 295201 arXiv:0803.0785 [hep-th]
- [33] Nagatomo K and Tsuchiya A 2011 *Adv. Stud. Pure Math.* **61** 1–49 arXiv:0902.4607 [math.QA]
- [34] Tsuchiya A and Wood S 2013 *J. Phys.* **A46** 445203 arXiv:1201.0419 [hep-th]
- [35] Gainutdinov A, Saleur H and Tipunin I Y Lattice W-algebras and logarithmic CFTs arXiv:1212.1378 [hep-th]

- [36] Feigin B, Gainutdinov A, Semikhatov A and Tipunin I Y 2006 *Nucl. Phys.* **B757** 303–343 arXiv:hep-th/0606196
- [37] Rasmussen J 2009 *Nucl. Phys.* **B807** 495–533 arXiv:0805.2991 [hep-th]
- [38] Gaberdiel M, Runkel I and Wood S 2009 *J. Phys.* **A42** 325403 arXiv:0905.0916 [hep-th]
- [39] Wood S 2010 *J. Phys.* **A43** 045212 arXiv:0907.4421 [hep-th]
- [40] Adamović D and Milas A 2010 *Int. Math. Res. Not.* **2010** 3896–3934 arXiv:0908.4053 [math.QA]
- [41] Gaberdiel M, Runkel I and Wood S 2011 *J. Phys.* **A44** 015204 arXiv:1008.0082 [hep-th]
- [42] Adamović D and Milas A 2012 *J. Math. Phys.* **53** 073511 arXiv:1202.6667 [math.QA]
- [43] Tsuchiya A and Wood S 2014 *Int. Math. Res. Not.* (to appear) arXiv:1302.6435 [hep-th]
- [44] Semikhatov A and Tipunin I Y 2013 *J. Phys.* **A46** 494011 arXiv:1301.2235 [math.QA]
- [45] Verlinde E 1988 *Nucl. Phys.* **B300** 360–376
- [46] Flohr M 1996 *Int. J. Mod. Phys.* **A11** 4147–4172 arXiv:hep-th/9509166
- [47] Miyamoto M 2004 *Duke Math. J.* **122** 51–91 arXiv:math/0209101 [math.QA]
- [48] Fuchs J, Schweigert C and Stigner C 2013 *J. Phys.* **A46** 494012 arXiv:1302.4683 [hep-th]
- [49] Fuchs J, Hwang S, Semikhatov A and Tipunin I Y 2004 *Comm. Math. Phys.* **247** 713–742 arXiv:hep-th/0306274
- [50] Flohr M and Knuth H On Verlinde-Like Formulas in $c(p, 1)$ Logarithmic Conformal Field Theories arXiv:0705.0545 [math-ph]
- [51] Gainutdinov A and Tipunin I Y 2009 *J. Phys.* **A42** 315207 arXiv:0711.3430 [hep-th]
- [52] Pearce P, Rasmussen J and Ruelle P 2010 *J. Phys.* **A43** 045211 arXiv:0907.0134 [hep-th]
- [53] Ridout D and Wood S 2014 *Nucl. Phys.* **B880** 175–202 arXiv:1310.6479 [hep-th]
- [54] Kytölä K and Ridout D 2009 *J. Math. Phys.* **50** 123503 arXiv:0905.0108 [math-ph]
- [55] Creutzig T and Ridout D 2012 *Nucl. Phys.* **B865** 83–114 arXiv:1205.6513 [hep-th]
- [56] Creutzig T and Ridout D 2013 *Nucl. Phys.* **B875** 423–458 arXiv:1306.4388 [hep-th]
- [57] Creutzig T and Ridout D 2011 *Springer Proc. Math. Stat.* **36** 349–368 arXiv:1111.5049 [hep-th]
- [58] Babichenko A and Ridout D 2013 *J. Phys.* **A46** 125204 arXiv:1210.7094 [math-ph]
- [59] Morin-Duchesne A, Rasmussen J and Ridout D Kac Modules for the Logarithmic Minimal Models (in preparation)
- [60] Canagasabey M, Rasmussen J and Ridout D Fusion Rules and Kac Modules for the $N = 1$ Logarithmic Minimal Models (in preparation)
- [61] Schellekens A and Yankielowicz S 1990 *Int. J. Mod. Phys.* **A5** 2903–2952
- [62] Dong C, Li H and Mason G 1996 *Comm. Math. Phys.* **180** 671–707 arXiv:q-alg/9504008
- [63] Fuchs J, Schellekens A and Schweigert C 1996 *Nucl. Phys.* **B473** 323–366 arXiv:hep-th/9601078
- [64] Mathieu P and Ridout D 2007 *Nucl. Phys.* **B765** 201–239 arXiv:hep-th/0609226
- [65] Mathieu P and Ridout D 2007 *Nucl. Phys.* **B776** 365–404 arXiv:hep-th/0701250
- [66] Miyamoto M 1998 *J. Alg.* **201** 115–150 arXiv:hep-th/9612032
- [67] Lam C 2001 *J. Math. Soc. Japan* **53** 541–557
- [68] Huang Y Z and Lepowsky J 2013 *J. Phys.* **A46** 494009 arXiv:1304.7556 [hep-th]