

THE PRINCIPAL W-ALGEBRA OF $\mathfrak{psl}_{2|2}$

ZACHARY FEHILY, CHRISTOPHER RAYMOND AND DAVID RIDOUT

ABSTRACT. We study the structure and representation theory of the principal W -algebra W_{pr}^k of $V^k(\mathfrak{psl}_{2|2})$. The defining operator product expansions are computed, as is the Zhu algebra, and these results are used to classify irreducible highest-weight modules. In particular, for $k = \pm \frac{1}{2}$, W_{pr}^k is not simple and the corresponding simple quotient is the symplectic fermion vertex algebra. We use this fact, along with inverse hamiltonian reduction, to study relaxed highest-weight and logarithmic modules for the small $N = 4$ superconformal algebra at central charges -9 and -3 .

CONTENTS

1. Introduction	1
1.1. Outline and results	2
2. Computing the principal W -algebra	3
2.1. The Lie superalgebra $\mathfrak{psl}_{2 2}$	3
2.2. The principal reduction	4
2.3. Operator product expansions	6
3. Irreducible W_{pr}^k -modules	8
3.1. The principal Zhu algebra	8
3.2. Modules of the Zhu algebra	8
4. Inverse reduction and the $N = 4$ superconformal algebra	10
4.1. Inverse quantum hamiltonian reduction	10
4.2. Constructing weight $W_{\pm 1/2}^{\text{min}}$ -modules	12
4.3. Degenerations and spectral flows	13
4.4. Logarithmic $W_{\pm 1/2}^{\text{min}}$ -modules	15
References	18

1. INTRODUCTION

The simple Lie superalgebra $\mathfrak{psl}_{2|2}$ is somewhat unusual among the basic classical examples in that it has odd roots of multiplicity 2, which may be regarded as being both positive and negative. Nevertheless, it has received an enormous amount of attention due to its applications in physics. In particular, the “baby” version of the AdS/CFT correspondence on $\text{AdS}_3 \times S^3$ supersymmetrises to a sigma model on the Lie supergroup $\text{PSU}_{1,1|2}$ [12], leading to interest in Wess-Zumino-Witten models with $\widehat{\mathfrak{psl}}_{2|2}$ symmetry (see for example [22, 31, 33–35, 50]).

Intertwined with this story is that of the (small) $N = 4$ superconformal algebra. Introduced initially to describe spacetime supersymmetries of heterotic strings, its unitary representations were first studied in [24], motivated by applications to string theory on hyperKähler manifolds. More recently, there has been a resurgence in interest in $N = 4$ representation theory because of its role in such things as Mathieu moonshine and its variants [23]. However, this representation theory is still poorly understood in general. Classifications are only known for the unitary highest-weight modules [36, 37] and the irreducibles in the case of central charge $c = -9$ [1]. The $c = -9$ algebra is also the $p = 2$ member of the family of extensions $\mathcal{V}^{(p)}$ of $L_k(\mathfrak{sl}_2)$ at $k = -2 + \frac{1}{p}$, introduced in [1] and studied further in [5].

In a different direction, recent developments in the 3d/2d (and 4d/2d) correspondence has renewed interest in theories with $N = 4$ supersymmetry. In [14], it was argued that generic 3-dimensional gauge theories with an appropriate boundary condition support vertex-operator algebras on the boundary. The motivation here was that the data of the vertex-operator algebra, for example its fusion algebra, could be used to investigate a related topological quantum field theory (analogous to the well known story connecting Wess-Zumino-Witten models and 3d Chern–Simons theories).

A rich source of 3-dimensional gauge theories are the quiver gauge theories [21], in which the field content is determined by specifying a quiver. These theories have a rich associated geometry described by the corresponding quiver varieties of

Nakajima [44, 45]. One example, analysed in [10], is associated with the Jordan quiver for which the associated quiver variety is the Hilbert scheme $\text{Hilb}^n(\mathbb{C})$ of n -points in the complex plane. The authors use chiral quantisation to construct a vertex-operator algebra for each n that has $\text{Hilb}^n(\mathbb{C})$ as its associated variety. They also showed that for $n = 2$, their vertex-operator algebra is the small $N = 4$ superconformal algebra at $c = -9$ (tensored with some additional free-field vertex algebras). As such, a detailed understanding of $N = 4$ representation theory, for example its modular data and fusion rules, will shed light on the associated topological quantum field theory.

With this goal in mind, our interest is the relationship between the vertex superalgebras constructed from $\mathfrak{psl}_{2|2}$ and $N = 4$. In the formalism developed by Kac and Wakimoto, these are related by quantum hamiltonian reduction: the small $N = 4$ superconformal algebra is a minimal reduction of the affine vertex superalgebra constructed from $\mathfrak{psl}_{2|2}$ [38, 39]. But there is also a principal reduction which appears to have received little to no attention in the literature. This paper then amounts to the first steps towards understanding this reduction and its representation theory.

Given the recent interest in $N = 4$ representation theory, we aim to connect this to the representation theory of the principal reduction. Happily, the formalism of inverse quantum hamiltonian reduction is designed for this purpose. Originally introduced by Semikhatov [48] for the universal Virasoro and $\mathfrak{sl}_2$ vertex algebras, inverse reduction was extended to the simple quotients by Adamović in [2]. The latter also studied certain restriction functors that connected the representation theories, specifically sending irreducible highest-weight Virasoro modules to (spectral flows of) relaxed highest-weight $L_k(\mathfrak{sl}_2)$ -modules. Subsequent work generalised this to higher ranks and developed methods to prove that these $L_k(\mathfrak{sl}_2)$ -modules were generically irreducible [7] and that every irreducible relaxed $L_k(\mathfrak{sl}_2)$ -module could be obtained in this fashion [8]. Other recent works addressing inverse reduction include [3, 4, 25–30].

Inverse reduction therefore suggests a general path to understanding the representation theory of a nonrational simple affine vertex-operator algebra or W -algebra (associated to a simple Lie superalgebra $\mathfrak{g}$ at level k). One starts with the representation theory of the “exceptional” W -algebra of level k [11] and then use inverse reduction à la Adamović to reconstruct the representation theory of each successive “less reduced” W -algebra. One thereby ascends through the poset of level- k W -algebras corresponding to $\mathfrak{g}$ and k (parametrised by the even nilpotent orbits of $\mathfrak{g}$), iteratively building up their representation theories.

We remark that when $\mathfrak{g}$ is a Lie algebra and k is (co)admissible, the exceptional W -algebra is rational [11] and so its representation theory is well understood. This need not be the case if k is nonadmissible, see [8, §5.2], or $\mathfrak{g}$ is not a Lie algebra (or $\mathfrak{osp}_{1|2n}$), see [15]. In the case of interest here ($\mathfrak{g} = \mathfrak{psl}_{2|2}$), we identify the “exceptional” W -algebra for $k = \pm\frac{1}{2}$ as the symplectic fermions vertex-operator superalgebra [41]. Inverse reduction then allows us to investigate the representation theory of the small $N = 4$ superconformal algebra at central charge $c = -9$ (and also at $c = -3$).

1.1. Outline and results. We begin in Section 2 by recalling some basic facts about the Lie superalgebra $\mathfrak{psl}_{2|2}$. We then compute the principal W -algebra W_{pr}^k from the universal affine vertex-operator superalgebra $V^k(\mathfrak{psl}_{2|2})$ via quantum hamiltonian reduction. The operator product expansions that define W_{pr}^k are given in Theorem 2.1 and we record that $k = \pm\frac{1}{2}$ is a collapsing level with simple quotient isomorphic to symplectic fermions. We also conjecture the levels for which W_{pr}^k is not simple.

In Section 3, we study the lower-bounded weight modules of W_{pr}^k by computing the Zhu algebra $\text{Zhu}[W_{\text{pr}}^k]$ and determining the irreducible weight $\text{Zhu}[W_{\text{pr}}^k]$ -modules. Every irreducible lower-bounded W_{pr}^k -module is shown to be highest-weight and the dimensions of their top spaces are determined. In particular, the top spaces are always finite-dimensional. This confirms that the representation theory of W_{pr}^k is not too dissimilar to those of the principal W -algebras with $\mathfrak{g}$ a Lie algebra.

Section 4 then applies inverse hamiltonian reduction to W_{pr}^k . We first realise the minimal W -algebra W_{min}^k of $\mathfrak{psl}_{2|2}$ inside $W_{\text{pr}}^k \otimes \Pi$, where Π is the usual bosonisation of symplectic bosons. The argument determining when this realisation descends to the simple quotients W_k^{min} and W_k^{pr} is then sketched. Granting this, Section 4.2 restricts to the case $k = \pm\frac{1}{2}$ for which inverse reduction translates the well known representation theory of the symplectic fermions vertex-operator superalgebra into that of $W_{\pm\frac{1}{2}}^{\text{min}}$. We thereby recover the classification of irreducible highest-weight modules [1] and extend it to irreducible relaxed highest-weight modules (in both the Neveu–Schwarz and Ramond sectors). The (super)characters of the latter are easily computed, though we leave the potentially subtle analysis of their modularity to future work.

In anticipation of this modularity study, we turn to the degenerations and spectral flows of the relaxed families in Section 4.3. Interestingly, each member of these families is completely reducible, decomposing as a direct sum of two generically irreducible relaxed highest-weight modules. We also exhibit the structures of the reducible examples using Loewy diagrams. Here, we observe a qualitative difference between the $k = -\frac{1}{2}$ and $k = +\frac{1}{2}$ cases: unlike the former, one of the reducible relaxed modules of the latter has composition factors whose minimal conformal weights are strictly greater than that of the top space. We conclude in Section 4.4 by applying inverse reduction to the logarithmic module of symplectic fermions, thereby obtaining an uncountably infinite number of logarithmic $W_{\pm 1/2}^{\min}$ -modules. The structures of these modules are completely elucidated, even at degeneration points.

Acknowledgements. We thank Dražen Adamović, Thomas Creutzig, Justine Fasquel, Andy Linshaw and Sven Möller for helpful conversations on topics related to this research. CR's research is supported by the Deutsche Forschungsgemeinschaft through the Collaborative Research Centre 1624: Higher Structures, Moduli Spaces and Integrability, project number 50663264. DR's research is supported by the Australian Research Council Discovery Project DP210101502 and an Australian Research Council Future Fellowship FT200100431.

2. COMPUTING THE PRINCIPAL W-ALGEBRA

In this section, we introduce the principal W-algebras associated to the simple complex Lie superalgebra $\mathfrak{psl}_{2|2}$. For this, we follow the seminal work of Kac, Roan and Wakimoto [38] to identify generators and compute their defining operator product expansions explicitly.

2.1. The Lie superalgebra $\mathfrak{psl}_{2|2}$. Recall that $\mathfrak{sl}_{2|2}$ is the Lie superalgebra whose elements are the endomorphisms of the $\mathbb{Z}_2$ -graded vector superspace $\mathbb{C}^{2|2}$ with zero supertrace. The quotient of this superalgebra by the ideal spanned by the identity matrix is denoted by $\mathfrak{psl}_{2|2}$. Let E_{ij} , $i, j = 1, \dots, 4$, denote the 4×4 matrix whose (i, j) -th entry is 1 and whose other entries are 0. We fix a basis of $\mathfrak{psl}_{2|2}$ as follows, indicating parity and understanding that each E_{ij} is implicitly replaced by its image in the quotient:

$$(2.1) \quad \begin{aligned} E^1 = E_{12}, \quad H^1 = E_{11} - E_{22}, \quad F^1 = E_{21}, \quad E^2 = E_{34}, \quad H^2 = E_{33} - E_{44}, \quad F^2 = E_{43} & \quad (\text{even}), \\ e^{++} = +E_{14}, \quad e^{+-} = -E_{13}, \quad e^{-+} = +E_{24}, \quad e^{--} = -E_{23}, & \\ f^{++} = -E_{32}, \quad f^{+-} = -E_{42}, \quad f^{-+} = +E_{31}, \quad f^{--} = +E_{41} & \quad (\text{odd}). \end{aligned}$$

Here are some useful facts about $\mathfrak{psl}_{2|2}$ and the basis that we have chosen:

- The even subalgebra is isomorphic to $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$. Both $\{E^1, H^1, F^1\}$ and $\{E^2, H^2, F^2\}$ are $\mathfrak{sl}_2$ -bases.
- The four odd elements labelled with an e span a copy of the tensor product of two fundamental $\mathfrak{sl}_2$ -modules. The signs in (2.1) ensure that the actions of the E^i and F^i , $i = 1, 2$, just exchange the i -th $\pm$ index. For example,

$$(2.2) \quad F^1 e^{++} = e^{-+} \quad \text{and} \quad E^2 e^{+-} = e^{++}.$$

- The previous property also holds for the odd elements labelled with an f .
- The odd roots have multiplicity 2. The corresponding root vectors are nilpotent: $[e^{\pm\pm}, e^{\pm\pm}] = [f^{\pm\pm}, f^{\pm\pm}] = 0$. (Here and throughout, $[\cdot, \cdot]$ denotes a Lie superbracket, modelled on the anticommutator if both elements are odd and on the commutator if both elements are even or one is even and one odd.)
- In fact, the odd elements labelled by an e all anticommute with one another. The same is true for the odd elements labelled by an f .
- The nonzero anticommutators are as follows:

$$(2.3) \quad \begin{aligned} [e^{++}, f^{+-}] &= -E^1, & [e^{++}, f^{++}] &= +E^2, & [e^{++}, f^{--}] &= +\frac{1}{2}(H^1 - H^2), \\ [e^{+-}, f^{++}] &= +E^1, & [e^{+-}, f^{+-}] &= -\frac{1}{2}(H^1 + H^2), & [e^{+-}, f^{--}] &= -F^2, \\ [e^{-+}, f^{++}] &= -E^2, & [e^{-+}, f^{+-}] &= \frac{1}{2}(H^1 + H^2), & [e^{-+}, f^{--}] &= +F^1, \\ [e^{--}, f^{++}] &= -\frac{1}{2}(H^1 - H^2), & [e^{--}, f^{+-}] &= +F^2, & [e^{--}, f^{++}] &= -F^1. \end{aligned}$$




FIGURE 1. The roots of the simple Lie superalgebra $\mathfrak{psI}_{2|2}$, labelled by their root vectors (the odd roots have multiplicity 2). The horizontal axis indicates the eigenvalue under the adjoint action of H^1 while the vertical axis records it for H^2 .

- The Killing form is identically zero, but the supertrace form in the defining representation (2.1) is supersymmetric and nondegenerate. The nonzero entries of this supertrace form are as follows:

$$(2.4) \quad \begin{aligned} \langle E^1, F^1 \rangle &= +1, & \langle H^1, H^1 \rangle &= +2, & \langle E^2, F^2 \rangle &= -1, & \langle H^2, H^2 \rangle &= -2, \\ \langle e^{++}, f^{--} \rangle &= +1, & \langle e^{+-}, f^{+-} \rangle &= -1, & \langle e^{-+}, f^{+-} \rangle &= -1, & \langle e^{--}, f^{++} \rangle &= +1. \end{aligned}$$

We picture the root system of $\mathfrak{psI}_{2|2}$ in Figure 1, indicating by position the eigenvalues with respect to the basis $\{H^1, H^2\}$ of the Cartan subalgebra.

The Weyl group of $\mathfrak{psI}_{2|2}$ is, by definition, that of the even subalgebra $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$, hence it is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$. However, the automorphism group of the root system is clearly isomorphic to D_4 . The additional automorphisms are generated by the reflection that fixes the odd root of weight $(1, 1)$. This automorphism corresponds to swapping the two copies of $\mathfrak{sl}_2$. It moreover lifts to an order-2 automorphism ω of $\mathfrak{psI}_{2|2}$ defined by

$$(2.5) \quad E^1 \leftrightarrow E^2, \quad H^1 \leftrightarrow H^2, \quad F^1 \leftrightarrow F^2, \quad e^{++} \leftrightarrow f^{++}, \quad e^{+-} \leftrightarrow f^{+-}, \quad e^{-+} \leftrightarrow f^{+-}, \quad e^{--} \leftrightarrow f^{--}.$$

Interestingly, this automorphism does not leave the supertrace form invariant, but instead negates it:

$$(2.6) \quad \langle \omega(A), \omega(B) \rangle = -\langle A, B \rangle, \quad A, B \in \mathfrak{psI}_{2|2}.$$

2.2. The principal reduction. The (untwisted) affine Kac–Moody superalgebra $\widehat{\mathfrak{psI}}_{2|2}$ is spanned by elements A_n and K , for $A \in \mathfrak{psI}_{2|2}$ and $n \in \mathbb{Z}$, where K is even and A_n inherits its parity from that of A . As usual, K is central and the Lie superbrackets for the A_n are given by

$$(2.7) \quad [A_m, B_n] = [A, B]_{m+n} + m\langle A, B \rangle \delta_{m+n,0} K, \quad A, B \in \mathfrak{psI}_{2|2}, \quad m, n \in \mathbb{Z}.$$

Because of (2.6), the automorphism ω of $\mathfrak{psI}_{2|2}$ lifts to an automorphism of $\widehat{\mathfrak{psI}}_{2|2}$ by defining

$$(2.8) \quad \widehat{\omega}(A_n) = \omega(A)_n \quad \text{and} \quad \widehat{\omega}(K) = -K, \quad A \in \mathfrak{psI}_{2|2}, \quad n \in \mathbb{Z}.$$

The universal level- k affine vertex superalgebra $V^k(\mathfrak{psI}_{2|2})$ is defined, as always, to be strongly and freely generated by the fields $A(z) = \sum_{n \in \mathbb{Z}} A_n z^{-n-1}$, $A \in \mathfrak{psI}_{2|2}$, subject to the operator product expansions

$$(2.9) \quad A(z)B(w) \sim \frac{\langle A, B \rangle k \mathbb{1}}{(z-w)^2} + \frac{[A, B](z)}{z-w}, \quad A, B \in \mathfrak{psI}_{2|2}.$$

Here, $\mathbb{1}$ denotes the identity field and the A_n furnish a representation of $\widehat{\mathfrak{psI}}_{2|2}$ on $V^k(\mathfrak{psI}_{2|2})$ in which K acts as multiplication by the level $k \in \mathbb{C}$. We note that the automorphism $\widehat{\omega}$ of $\widehat{\mathfrak{psI}}_{2|2}$ again lifts, but this time to an *isomorphism* between $V^k(\mathfrak{psI}_{2|2})$ and $V^{-k}(\mathfrak{psI}_{2|2})$. It follows, in particular, that the representation theory of this vertex superalgebra (and any of its quotients) is independent of the sign of the level.

Because the supertrace form (2.4) is nondegenerate, the Sugawara construction equips $V^k(\mathfrak{psI}_{2|2})$ with the structure of a vertex-operator superalgebra, when the level is noncritical. The dual Coxeter number of $\mathfrak{psI}_{2|2}$ is 0 because the Killing form vanishes, so for all $k \neq 0$, the energy-momentum tensor takes the form

$$(2.10) \quad T = \frac{1}{2k} \left(\frac{1}{2} :H^1 H^1: + :E^1 F^1: + :F^1 E^1: - \frac{1}{2} :H^2 H^2: - :E^2 F^2: - :F^2 E^2: \right. \\ \left. - (:e^{++} f^{--}: - :f^{--} e^{++}: - :e^{+-} f^{-+}: + :f^{-+} e^{+-}: - :e^{-+} f^{+-}: + :f^{+-} e^{-+}: + :e^{--} f^{++}: - :f^{++} e^{--}:) \right).$$

The central charge is the superdimension of $\mathfrak{psl}_{2|2}$, namely -2 .

As there are two nilpotent orbits in $\mathfrak{sl}_2$, there are four even orbits in $\mathfrak{psl}_{2|2}$. Along with the zero orbit, there are two minimal orbits, for which F^1 and F^2 are representatives, and the principal one, for which $F^1 + F^2$ is a representative. It is well known that quantum hamiltonian reduction $W_{\min}^k$ with respect to one of the minimal nilpotents is isomorphic to the universal (small) $N = 4$ superconformal vertex-operator superalgebra of central charge $-6(k+1)$, see [38, Rem. 4.1] and [39, §8.4]. Applying $\widehat{\omega}$, we see that the other minimal reduction is isomorphic to $W_{\min}^{-k}$ with central charge $6(k-1)$. We leave further consideration of these minimal reductions to Section 4.

Our interest is in the principal reduction W_{pr}^k , which does not seem to have received as much attention. We observe that $\{E^1 + E^2, H^1 + H^2, F^1 + F^2\}$ is an $\mathfrak{sl}_2$ -triple and that the adjoint action of $H^1 + H^2$ gives a good even grading on $\mathfrak{psl}_{2|2}$:

$$(2.11) \quad \mathfrak{psl}_{2|2} = \mathfrak{psl}_{2|2}^{(-2)} \oplus \mathfrak{psl}_{2|2}^{(0)} \oplus \mathfrak{psl}_{2|2}^{(+2)}, \quad \begin{aligned} \mathfrak{psl}_{2|2}^{(+2)} &= \text{span}\{E^1, E^2 \mid e^{++}, f^{++}\}, \\ \mathfrak{psl}_{2|2}^{(0)} &= \text{span}\{H^1, H^2 \mid e^{+-}, e^{-+}, f^{+-}, f^{-+}\}, \\ \mathfrak{psl}_{2|2}^{(-2)} &= \text{span}\{F^1, F^2 \mid e^{--}, f^{--}\}. \end{aligned}$$

By [39, Thm. 4.1], W_{pr}^k is then strongly and freely generated, up to some operator product expansions that must be computed, by two even fields of conformal weight 2 and four odd fields of conformal weights 1, 1, 2 and 2.

Following the recipe in [39], we associate to each even basis element of $\mathfrak{psl}_{2|2}^{(+2)}$ a fermionic ghost vertex superalgebra FG^i , $i = 1, 2$, and to each odd basis element a bosonic ghost vertex algebra BG^j , $j = e, f$. These are strongly and freely generated by fields satisfying

$$(2.12) \quad b^i(z)c^i(w) \sim \frac{\mathbb{1}}{z-w} \quad \text{and} \quad \beta^j(z)\gamma^j(w) \sim -\frac{\mathbb{1}}{z-w},$$

respectively, with all other operator product expansions of the generators being regular. We next form the graded complex

$$(2.13) \quad C = V^k(\mathfrak{psl}_{2|2}) \otimes FG^1 \otimes FG^2 \otimes BG^e \otimes BG^f,$$

in which the b^i and β^j have grade -1 , the c^i and γ^j have grade $+1$, and all other generators (and vacuum states) have grade 0. This becomes a differential graded complex upon taking the differential to be the zero mode of

$$(2.14) \quad D = (E^1 + \mathbb{1})c^1 + (E^2 - \mathbb{1})c^2 - e^{++}\gamma^e - f^{++}\gamma^f.$$

D_0 is odd, has grade $+1$ and is easily checked to square to zero. As a vertex superalgebra, W_{pr}^k is the grade-0 cohomology of C with respect to D_0 .

To give W_{pr}^k , $k \neq 0$, the structure of a vertex-operator superalgebra, we choose an energy-momentum tensor L on C . This needs to make D homogeneous of conformal weight 1, so we take

$$(2.15) \quad L = T + \frac{1}{2}(\partial H^1 + \partial H^2) + :\partial b^1 c^1: + :\partial b^2 c^2: + :\partial \beta^e \gamma^e: + :\partial \beta^f \gamma^f:.$$

It is easy to check that C is now a vertex-operator superalgebra of central charge -2 and that $D_0 L = 0$. Moreover, L defines an energy-momentum tensor on W_{pr}^k , as desired, by [38, Thm. 2.2].

The next step is to construct the “parenthetical building blocks” of the generators of W_{pr}^k . These are obtained by correcting some of the generators of $V^k(\mathfrak{psl}_{2|2})$ by adding bilinear terms in the ghost fields, see [39, Eq. (2.4)]. We need to compute them for all elements in $\mathfrak{psl}_{2|2}^{(0)}$, the results for our basis being

$$(2.16) \quad \begin{aligned} H^{(1)} &= H^1 + 2:b^1 c^1: + :\beta^e \gamma^e: + :\beta^f \gamma^f:, & e^{(+)} &= e^{+-} - :\beta^e c^2: + :b^1 \gamma^f:, & e^{(-)} &= e^{-+} - :\beta^e c^1: - :b^2 \gamma^f:, \\ H^{(2)} &= H^2 + 2:b^2 c^2: + :\beta^e \gamma^e: + :\beta^f \gamma^f:, & f^{(+)} &= f^{+-} - :\beta^f c^2: - :b^1 \gamma^e:, & f^{(-)} &= f^{-+} - :\beta^f c^1: + :b^2 \gamma^e:. \end{aligned}$$

The generating fields of conformal weight 1 in W_{pr}^k are then (the images in cohomology of) the linear combinations of the parenthetical building blocks that commute with F [39, Thm. 4.1]. In this way, we find two odd generators:

$$(2.17) \quad \chi^e = e^{(+)} - e^{(-)} \quad \text{and} \quad \chi^f = f^{(+)} - f^{(-)}.$$

It is easy to check that these are D_0 -closed.

To determine the weight-2 generating fields of W_{pr}^k , we must find D_0 -closed corrections to the elements of $\mathfrak{psl}_{2|2}^{(-2)}$ (all of which commute with F). This time, [39] does not give a precise recipe, only guaranteeing that the corrections belong to the subalgebra generated by the parenthetical building blocks (2.16). Finding the corrections by brute force is quite tractable in this case (we used OPEDEFS [49]). The resulting four generators are

$$(2.18) \quad \begin{aligned} B^1 &= F^1 - \frac{1}{4} :H^{(1)}H^{(1)}: - \frac{k}{2} \partial H^{(1)} - :e^{(-+)}f^{(-+)}: - \frac{(2k-1)(3k+1)}{4k} : (e^{(+ -)} - e^{(-+)})(f^{(+ -)} - f^{(-+)}):, \\ B^2 &= F^2 + \frac{1}{4} :H^{(2)}H^{(2)}: - \frac{k}{2} \partial H^{(2)} - :e^{(+ -)}f^{(+ -)}: + \frac{(2k+1)(3k-1)}{4k} : (e^{(+ -)} - e^{(-+)})(f^{(+ -)} - f^{(-+)}):, \\ \psi^e &= e^{--} - \frac{1}{2} (:H^{(1)}e^{(+ -)}: - :H^{(2)}e^{(-+)}:) - \frac{2k-1}{4} \partial e^{(+ -)} - \frac{2k+1}{4} \partial e^{(-+)}, \\ \psi^f &= f^{--} - \frac{1}{2} (:H^{(1)}f^{(+ -)}: - :H^{(2)}f^{(-+)}:) - \frac{2k-1}{4} \partial f^{(+ -)} - \frac{2k+1}{4} \partial f^{(-+)}, \end{aligned}$$

where unique solutions were obtained by requiring in addition that the generators be primary with respect to L . We record that (up to D_0 -exact terms)

$$(2.19) \quad L = -\frac{1}{k}(B^1 + B^2) - \frac{1}{2k} : \chi^e \chi^f :.$$

2.3. Operator product expansions. Having explicitly computed the generating fields of the universal level- k principal W -algebra W_{pr}^k , we now turn to its operator product expansions. By [39, Thm. 4.1 and Rem. 4.2], W_{pr}^k admits a Poincaré–Birkhoff–Witt basis in the modes of the generating fields. It follows that W_{pr}^k is strongly and freely generated by B^1 , B^2 , χ^e , χ^f , ψ^e and ψ^f , subject only to their operator product expansions. In principle, these relations might only close modulo D_0 -exact fields. Happily, this is not the case.

The resulting operator product expansions are nevertheless somewhat messy. To bring out some of the structural features, we first present those of the generating fields of conformal weight 1:

$$(2.20) \quad \chi^e(z)\chi^e(w) \sim 0, \quad \chi^e(z)\chi^f(w) \sim \frac{2k\mathbb{1}}{(z-w)^2}, \quad \chi^f(z)\chi^f(w) \sim 0.$$

The weight-1 fields of W_{pr}^k thus generate a subalgebra isomorphic to a pair of symplectic fermions [41]. We may even remove the apparent level dependence by defining renormalised symplectic fermions thusly:

$$(2.21) \quad \chi = \frac{1}{\sqrt{k}} \chi^e \quad \text{and} \quad \bar{\chi} = \frac{1}{\sqrt{k}} \chi^f.$$

To choose a better basis of weight-2 generating fields, we require that they transform “cleanly” under the action of the symplectic fermion subalgebra. More precisely, the space of weight-2 generators forms a representation of the zero modes of the symplectic fermions vertex-operator superalgebra $\text{SF} = \mathbb{V}^k(\mathfrak{psl}_{1|1})$ (these are isomorphic for all $k \neq 0$), so we can look for a basis that respects the structure of this representation. Since the zero modes of the generators define a copy of $\mathfrak{psl}_{1|1}$, a Lie superalgebra with relatively few representations, it is in fact rather easy to identify ours. Indeed, if we define

$$(2.22) \quad S = -\frac{1}{k}(B^1 + B^2), \quad H = \frac{1}{2}(B^2 - B^1) - \frac{1}{2}S, \quad \psi = \frac{1}{\sqrt{k}}\psi^e \quad \text{and} \quad \bar{\psi} = \frac{1}{\sqrt{k}}\psi^f,$$

it is straightforward to verify (again, using OPEDEFS) that

$$(2.23) \quad \chi_0 H = \psi, \quad \bar{\chi}_0 H = \bar{\psi}, \quad \chi_0 \bar{\psi} = +S, \quad \bar{\chi}_0 \psi = -S, \quad \text{and} \quad \chi_0 \psi = \bar{\chi}_0 \bar{\psi} = \chi_0 S = \bar{\chi}_0 S = 0.$$

The space spanned by the weight-2 generators is thus a copy of the unique projective module of $\mathfrak{psl}_{1|1}$, which is itself isomorphic to $U(\mathfrak{psl}_{1|1})$.

Note that the energy-momentum tensor of W_{pr}^k is now given by

$$(2.24) \quad L = S - \frac{1}{2} : \chi \bar{\chi} :.$$

Because of this, we may replace S by L in our set of strong generators. We record the defining operator product expansions of W_{pr}^k in the following theorem.

Theorem 2.1. *The principal quantum hamiltonian reduction W_{pr}^k of $V^k(\mathfrak{psl}_{2|2})$ is strongly and freely generated by two even elements, L and H , and four odd elements, χ , $\bar{\chi}$, ψ and $\bar{\psi}$. Here, L is a conformal vector of central charge -2 and the generators H , χ , $\bar{\chi}$, ψ and $\bar{\psi}$ are Virasoro primaries with respective conformal weights 2 , 1 , 1 , 2 and 2 . The remaining operator product expansions are as follows:*

$$\begin{aligned}
 (2.25a) \quad & \chi(z)\chi(w) \sim \bar{\chi}(z)\bar{\chi}(w) \sim 0, \quad \chi(z)\bar{\chi}(w) \sim \frac{2\mathbb{1}}{(z-w)^2}, \\
 (2.25b) \quad & \chi(z)\psi(w) \sim \bar{\chi}(z)\bar{\psi}(w) \sim 0, \quad \chi(z)\bar{\psi}(w) \sim +\frac{S(w)}{z-w}, \quad \bar{\chi}(z)\psi(w) \sim -\frac{S(w)}{z-w}, \\
 (2.25c) \quad & \chi(z)H(w) \sim -3\left(k^2 - \frac{1}{4}\right)\frac{\chi(w)}{(z-w)^2} + \frac{\psi(w)}{z-w}, \quad \bar{\chi}(z)H(w) \sim -3\left(k^2 - \frac{1}{4}\right)\frac{\bar{\chi}(w)}{(z-w)^2} + \frac{\bar{\psi}(w)}{z-w}, \\
 & H(z)\psi(w) \sim \frac{3}{2}\left(k^2 - \frac{1}{4}\right)\left[\frac{\chi(w)}{(z-w)^3} + \frac{\frac{1}{2}\partial\chi(w)}{(z-w)^2} + \frac{\frac{1}{6}\partial^2\chi(w)}{z-w}\right] + \frac{\frac{1}{4}\partial\psi(w) - :\chi(H + \frac{3k^2}{2}S): (w)}{z-w}, \\
 (2.25d) \quad & H(z)\bar{\psi}(w) \sim \frac{3}{2}\left(k^2 - \frac{1}{4}\right)\left[\frac{\bar{\chi}(w)}{(z-w)^3} + \frac{\frac{1}{2}\partial\bar{\chi}(w)}{(z-w)^2} + \frac{\frac{1}{6}\partial^2\bar{\chi}(w)}{z-w}\right] + \frac{\frac{1}{4}\partial\bar{\psi}(w) - :\bar{\chi}(H + \frac{3k^2}{2}S): (w)}{z-w}, \\
 & H(z)H(w) \sim -3(3k^2 - 1)\left(k^2 - \frac{1}{4}\right)\left[\frac{\mathbb{1}}{(z-w)^4} + \frac{:\chi\bar{\chi}: (w)}{(z-w)^2} + \frac{\frac{1}{2}\partial:\chi\bar{\chi}: (w)}{z-w}\right] + \frac{7k^2 - 1}{2}\left[\frac{S(w)}{(z-w)^2} + \frac{\frac{1}{2}\partial S(w)}{z-w}\right], \\
 (2.25e) \quad & \psi(z)\psi(w) \sim \frac{:\chi\psi: (w)}{z-w}, \quad \psi(z)\bar{\psi}(w) \sim 3\left(k^2 - \frac{1}{4}\right)\left[\frac{\mathbb{1}}{(z-w)^4} + \frac{:\chi\bar{\chi}: (w)}{(z-w)^2} + \frac{\frac{1}{2}\partial:\chi\bar{\chi}: (w)}{z-w}\right] \\
 & \bar{\psi}(z)\bar{\psi}(w) \sim \frac{:\bar{\chi}\bar{\psi}: (w)}{z-w}, \quad -\left[\frac{2H(w) + \frac{1}{2}S(w)}{(z-w)^2} + \frac{\partial H(w) + \frac{1}{4}\partial S(w) - \frac{1}{2}:\chi\bar{\psi}: (w) - \frac{1}{2}:\bar{\chi}\psi: (w)}{z-w}\right].
 \end{aligned}$$

In this theorem, we chose to present the defining operator product expansions (2.25) so that the right-hand sides involve S rather than L , in order to respect the $\mathfrak{psl}_{1|1}$ -symmetry. For completeness, we also give the operator product expansions in which the left-hand sides involve S :

$$\begin{aligned}
 (2.26a) \quad & \chi(z)S(w) \sim \bar{\chi}(z)S(w) \sim 0, \quad S(z)S(w) \sim \frac{2S(w)}{(z-w)^2} + \frac{\partial S(w)}{z-w}, \\
 (2.26b) \quad & S(z)\psi(w) \sim \frac{2\psi(w)}{(z-w)^2} + \frac{\partial\psi(w) - \frac{1}{2}:\chi S: (w)}{z-w}, \quad S(z)\bar{\psi}(w) \sim \frac{2\bar{\psi}(w)}{(z-w)^2} + \frac{\partial\bar{\psi}(w) - \frac{1}{2}:\bar{\chi} S: (w)}{z-w}, \\
 (2.26c) \quad & S(z)H(w) \sim -3\left(k^2 - \frac{1}{4}\right)\left[\frac{\mathbb{1}}{(z-w)^4} + \frac{:\chi\bar{\chi}: (w)}{(z-w)^2} + \frac{\frac{1}{2}\partial:\chi\bar{\chi}: (w)}{z-w}\right] \\
 & \quad + \frac{2H(w) + \frac{1}{2}S(w)}{(z-w)^2} + \frac{\partial H(w) + \frac{1}{2}:\chi\bar{\psi}: (w) - \frac{1}{2}:\bar{\chi}\psi: (w)}{z-w}.
 \end{aligned}$$

We record the following observations about W_{pr}^k :

- The operator product expansions (2.25) only depend on k^2 , not k , so that $W_{\text{pr}}^k \cong W_{\text{pr}}^{-k}$. This is an obvious consequence of the isomorphism $V^k(\mathfrak{psl}_{2|2}) \cong V^{-k}(\mathfrak{psl}_{2|2})$, noted in Section 2.2, and the fact that the principal nilpotent $F^1 + F^2 \in \mathfrak{psl}_{2|2}$ is invariant under ω .
- S is an energy-momentum tensor of central charge 0. It commutes with the symplectic fermions subalgebra generated by χ and $\bar{\chi}$. It would be interesting to identify the commutant of this subalgebra in W_{pr}^k .
- When $k^2 = \frac{1}{4}$ (hence $k = \pm\frac{1}{2}$), the level is “collapsing” [6], meaning that the simple quotient $W_{\text{pr}}^{\text{pr}}$ of W_{pr}^k is strongly generated by just its weight-1 fields. To see this, note that H , S , ψ and $\bar{\psi}$ generate an ideal of $W_{\text{pr}}^{\pm 1/2}$ and that the quotient is the symplectic fermions subalgebra (which is simple): $W_{\pm 1/2}^{\text{pr}} \cong \text{SF}$.
- W_{pr}^k admits a horizontal grading in which χ and ψ have grade +1, $\bar{\chi}$ and $\bar{\psi}$ have grade −1, and H , S and L have grade 0. Its graded character (ch^+) and supercharacter (ch^-) are thus

$$(2.27) \quad \text{ch}^\pm[W_{\text{pr}}^k](y; q) = q^{1/12} \prod_{i=1}^{\infty} \frac{(1 \pm yq^i)(1 \pm y^{-1}q^i)(1 \pm yq^{i+1})(1 \pm y^{-1}q^{i+1})}{(1 - q^{i+1})^2}.$$

- Numerical investigations with the analogue of the Shapovalov form on W_{pr}^k show that there are singular vectors of conformal weight 2, if $k = \pm \frac{1}{2}$, weight 4 if $k = \pm \frac{1}{3}, \pm \frac{3}{2}$, and weight 6, if $k = \pm \frac{1}{4}, \pm \frac{2}{3}, \pm \frac{5}{2}$. We therefore conjecture that W_{pr}^k is a simple vertex-operator superalgebra unless $k \in \mathbb{Q} \setminus \mathbb{Z}$. We also conjecture that when W_{pr}^k is not simple, then the singular vector of minimal conformal weight has grade +1 and multiplicity 1.

3. IRREDUCIBLE W_{pr}^k -MODULES

We next study the representation theory of W_{pr}^k . Being a principal W -algebra, we expect that its irreducible modules are all ordinary, meaning that the L_0 -eigenspaces are all finite-dimensional. This is proven under the additional hypothesis that the irreducible is lower bounded. We also conjecture that every irreducible W_{pr}^k -module is lower bounded.

3.1. The principal Zhu algebra. The Zhu algebra $\text{Zhu}[V]$ of a vertex superalgebra V is the unital associative superalgebra of zero modes of the fields, restricted to act on states that are annihilated by all positive modes. Introduced formally in [40, 51], $\text{Zhu}[V]$ may be realised as a quotient of V with an associative product given by

$$(3.1) \quad A \circ B = \sum_{j=0}^{\Delta_A} \binom{\Delta_A}{j} (A_{-\Delta_A+j} B)_{\circ}, \quad A, B \in V.$$

Here, $A_{\circ}$ denotes the image of A in $\text{Zhu}[V]$ and Δ_A is the conformal weight of A (which is assumed to be homogeneous). It inherits its parity from that of A . The quotient operation will not be important for us here, except to note that it includes the identification

$$(3.2) \quad (\partial A)_{\circ} = -\Delta_A A_{\circ}, \quad A \in V.$$

It is now easy to use (3.1) and (3.2) to deduce relations in $\text{Zhu}[W_{\text{pr}}^k]$ involving the images of the strong generators:

$$(3.3) \quad \begin{aligned} (\chi_{\circ})^2 &= (\bar{\chi}_{\circ})^2 = 0, & [\chi_{\circ}, S_{\circ}] &= [\bar{\chi}_{\circ}, S_{\circ}] = 0, & [\chi_{\circ}, H_{\circ}] &= \psi_{\circ}, & [\bar{\chi}_{\circ}, H_{\circ}] &= \bar{\psi}_{\circ}, \\ [\chi_{\circ}, \bar{\chi}_{\circ}] &= 0, & [\chi_{\circ}, \psi_{\circ}] &= [\bar{\chi}_{\circ}, \bar{\psi}_{\circ}] = 0, & [\chi_{\circ}, \bar{\psi}_{\circ}] &= +S_{\circ}, & [\bar{\chi}_{\circ}, \psi_{\circ}] &= -S_{\circ}, \\ (\psi_{\circ})^2 &= \frac{1}{2} \chi_{\circ} \psi_{\circ}, & (\bar{\psi}_{\circ})^2 &= \frac{1}{2} \bar{\chi}_{\circ} \bar{\psi}_{\circ}, & [\psi_{\circ}, S_{\circ}] &= \frac{1}{2} \chi_{\circ} S_{\circ}, & [\bar{\psi}_{\circ}, S_{\circ}] &= \frac{1}{2} \bar{\chi}_{\circ} S_{\circ}, \\ [\psi_{\circ}, \bar{\psi}_{\circ}] &= \frac{1}{2} (\chi_{\circ} \bar{\psi}_{\circ} + \bar{\chi}_{\circ} \psi_{\circ}), & [S_{\circ}, H_{\circ}] &= \frac{1}{2} (\chi_{\circ} \bar{\psi}_{\circ} + \psi_{\circ} \bar{\chi}_{\circ}), \\ [\psi_{\circ}, H_{\circ}] &= \chi_{\circ} (H_{\circ} + \frac{3k^2}{2} L_{\circ} + \frac{1}{4} (k^2 - \frac{1}{4}) \mathbb{1}_{\circ}) - \frac{1}{2} \psi_{\circ}, & [\bar{\psi}_{\circ}, H_{\circ}] &= \bar{\chi}_{\circ} (H_{\circ} + \frac{3k^2}{2} L_{\circ} + \frac{1}{4} (k^2 - \frac{1}{4}) \mathbb{1}_{\circ}) - \frac{1}{2} \bar{\psi}_{\circ}. \end{aligned}$$

We shall not do so here, but one can prove that the images of the strong generators, along with the relations (3.3), comprise a presentation of $\text{Zhu}[W_{\text{pr}}^k]$. Instead, we record some useful observations:

- As usual, $L_{\circ} = S_{\circ} + \frac{1}{2} \bar{\chi}_{\circ} \chi_{\circ}$ is central.
- The Zhu-images of the odd weight-2 fields are not nilpotent of order 2, as one might expect, but of order 3:

$$(3.4) \quad (\psi_{\circ})^3 = \frac{1}{2} \chi_{\circ} (\psi_{\circ})^2 = \frac{1}{4} (\chi_{\circ})^2 \psi_{\circ} = 0.$$

- The horizontal grading of W_{pr}^k descends to a grading of $\text{Zhu}[W_{\text{pr}}^k]$ in which $\chi_{\circ}$ and $\psi_{\circ}$ have grade +1, $\bar{\chi}_{\circ}$ and $\bar{\psi}_{\circ}$ have grade -1, and $H_{\circ}$, $L_{\circ}$ and $S_{\circ}$ have grade 0.
- A straightforward consequence of the relations (3.3) is that monomials in the generators of $\text{Zhu}[W_{\text{pr}}^k]$ may be consistently ordered so that the odd elements with bars (no bars) are placed to the left (right) of the even elements. For example, the juxtaposition $\psi_{\circ} H_{\circ}$ may be so reordered by writing

$$(3.5) \quad \psi_{\circ} H_{\circ} = H_{\circ} \psi_{\circ} + \chi_{\circ} (H_{\circ} + \frac{3k^2}{2} L_{\circ} + \frac{1}{4} (k^2 - \frac{1}{4}) \mathbb{1}_{\circ}) - \frac{1}{2} \psi_{\circ} = (H_{\circ} + \frac{1}{2} \mathbb{1}_{\circ}) \psi_{\circ} + (H_{\circ} + \frac{3k^2}{2} L_{\circ} + \frac{1}{4} (k^2 - \frac{1}{4}) \mathbb{1}_{\circ}) \chi_{\circ}.$$

This Poincaré–Birkhoff–Witt-style ordering suggests that $\text{Zhu}[W_{\text{pr}}^k]$ admits a well behaved highest-weight theory.

3.2. Modules of the Zhu algebra. Let us define a weight vector in a $\text{Zhu}[W_{\text{pr}}^k]$ -module to be a simultaneous eigenvector of $H_{\circ}$ and $L_{\circ}$. As usual, weight vectors always exist in (nonzero) finite-dimensional modules. We will call a weight vector a highest-weight vector if it is annihilated by both $\chi_{\circ}$ and $\psi_{\circ}$. Obviously, a highest-weight vector is also an eigenvector of $S_{\circ}$ (and the eigenvalue matches that of $L_{\circ}$).

Lemma 3.1. *If a $\text{Zhu}[W_{\text{pr}}^k]$ -module has a weight vector, then it has a highest-weight vector.*

Proof. Let v denote the hypothesised weight vector, so that $L_\circ v = \Delta v$ and $H_\circ v = hv$, for some $\Delta, h \in \mathbb{C}$. Then:

- If $\chi_\circ v = \psi_\circ v = 0$, then v is already highest-weight.
- If $\psi_\circ v = 0$ but $\chi_\circ v \neq 0$, then $\chi_\circ v$ is weight because $L_\circ$ is central and $H_\circ \chi_\circ v = \chi_\circ H_\circ v - \psi_\circ v = h\chi_\circ v$. But, $\chi_\circ \chi_\circ v = 0$ and $\psi_\circ \chi_\circ v = -\chi_\circ \psi_\circ v = 0$, so $\chi_\circ v$ is highest-weight.
- If $\chi_\circ v = 0$ but $\psi_\circ v \neq 0$, then $\psi_\circ v$ is weight because

$$(3.6) \quad H_\circ \psi_\circ v = \psi_\circ H_\circ v + \frac{1}{2} \psi_\circ v - \chi_\circ H_\circ v - \frac{3k^2}{2} \chi_\circ L_\circ v - \frac{1}{4} (k^2 - \frac{1}{4}) \chi_\circ v = (h + \frac{1}{2})v.$$

But, $\chi_\circ \psi_\circ v = -\psi_\circ \chi_\circ v = 0$ and $\psi_\circ \psi_\circ v = \frac{1}{2} \chi_\circ \psi_\circ v = 0$, so $\psi_\circ v$ is highest-weight.

- If $\chi_\circ v \neq 0$ and $\psi_\circ v \neq 0$ but $\chi_\circ \psi_\circ v = 0$, then

$$(3.7) \quad H_\circ \chi_\circ v = h\chi_\circ v - \psi_\circ v \quad \text{and} \quad H_\circ \psi_\circ v = (h + \frac{1}{2})\psi_\circ v - (h + \frac{3k^2}{2} \Delta + \frac{1}{4} (k^2 - \frac{1}{4})) \chi_\circ v.$$

Some linear combination of $\chi_\circ v$ and $\psi_\circ v$ is therefore a weight vector. Since $\chi_\circ \psi_\circ v = 0$, this linear combination is annihilated by both $\chi_\circ$ and $\psi_\circ$, hence it is a highest-weight vector.

- Finally, if $\chi_\circ \psi_\circ v \neq 0$ (so $\chi_\circ v \neq 0$ and $\psi_\circ v \neq 0$), then $H_\circ \chi_\circ \psi_\circ v = h\chi_\circ \psi_\circ v$ and $\chi_\circ \chi_\circ \psi_\circ v = \psi_\circ \chi_\circ \psi_\circ v = 0$, so $\chi_\circ \psi_\circ v$ is the desired highest-weight vector. ■

Inspection of (3.3) demonstrates that $H_\circ$, $L_\circ$, $\chi_\circ$ and $\psi_\circ$ generate a unital subalgebra of $\text{Zhu}[W_{\text{pr}}^k]$. We therefore may construct Verma modules in the usual way and realise all highest-weight modules, these being modules generated by a single highest-weight vector, as quotients. Let $\mathcal{V}_\circ^{h,\Delta}$ denote the Verma module generated by a highest-weight vector of $H_\circ$ -eigenvalue h and $L_\circ$ -eigenvalue Δ .

Proposition 3.2.

- $\mathcal{V}_\circ^{h,\Delta}$ is 4-dimensional, for all $h, \Delta \in \mathbb{C}$.
- $\mathcal{V}_\circ^{h,\Delta}$ is irreducible unless $\Delta = 0$.
- The irreducible quotient of $\mathcal{V}_\circ^{h,0}$ is 1-dimensional, for all $h \in \mathbb{C}$.

Proof. Let v be the generating highest-weight vector of $\mathcal{V}_\circ^{h,\Delta}$. Because we can order monomials so that the “negative” generators $\bar{\chi}_\circ$ and $\bar{\psi}_\circ$ act last, $\mathcal{V}_\circ^{h,\Delta}$ is spanned by the monomials obtained by acting on v with these negative generators. It is easy to check that these monomials are v , $\bar{\chi}_\circ v$, $\bar{\psi}_\circ v$ and $\bar{\chi}_\circ \bar{\psi}_\circ v$, hence that $\mathcal{V}_\circ^{h,\Delta}$ is 4-dimensional.

To investigate irreducibility, consider first the submodule of $\mathcal{V}_\circ^{h,\Delta}$ generated by the linear combination $w = a\bar{\chi}_\circ v + b\bar{\psi}_\circ v$, where $a, b \in \mathbb{C}$ are not both 0. Using (3.3), we find that $\chi_\circ w = \Delta b v$ and $\psi_\circ w = \Delta(-a + \frac{1}{2}b)v$. When $\Delta \neq 0$, it follows that this submodule contains v , for any $(a, b) \neq (0, 0)$, and is therefore $\mathcal{V}_\circ^{h,\Delta}$ itself. Similarly, $\Delta \neq 0$ implies that the submodule generated by $\bar{\chi}_\circ \bar{\psi}_\circ v$ contains $\chi_\circ \bar{\chi}_\circ \bar{\psi}_\circ v = -\Delta \bar{\chi}_\circ v$ and thus v , hence is also $\mathcal{V}_\circ^{h,\Delta}$. This proves the irreducibility of Verma modules when $\Delta \neq 0$.

When $\Delta = 0$, however, these calculations show that every nonzero linear combination of $\bar{\chi}_\circ v$ and $\bar{\psi}_\circ v$ is a highest-weight vector, hence generates a proper submodule. The sum of these proper submodules obviously contains $\bar{\chi}_\circ \bar{\psi}_\circ v$, so we conclude that the irreducible quotient of $\mathcal{V}_\circ^{h,0}$ is spanned by (the image of) v . ■

An irreducible $\text{Zhu}[W_{\text{pr}}^k]$ -module with a weight vector is thus a highest-weight module and so is either 4- or 1-dimensional. There are nevertheless reducible but indecomposable $\text{Zhu}[W_{\text{pr}}^k]$ -modules, also possessing weight vectors, of dimensions greater than 4. We shall not describe these here, but note instead that the action of $\text{Zhu}[W_{\text{pr}}^k]$ on Verma modules is not as nice as we might have hoped. In particular, the action of $S_\circ$ on $\mathcal{V}_\circ^{h,\Delta}$, $\Delta \neq 0$, is nondiagonalisable:

$$(3.8) \quad \begin{aligned} S_\circ v &= \Delta v, & S_\circ \bar{\chi}_\circ v &= \Delta \bar{\chi}_\circ v, \\ S_\circ \bar{\chi}_\circ \bar{\psi}_\circ v &= \Delta \bar{\chi}_\circ \bar{\psi}_\circ v, & S_\circ \bar{\psi}_\circ v &= -\frac{1}{2} \Delta \bar{\chi}_\circ v + \Delta \bar{\psi}_\circ v. \end{aligned}$$

The action of $H_\circ$ is even more opaque:

$$(3.9) \quad \begin{aligned} H_\circ v &= hv, & H_\circ \bar{\chi}_\circ v &= h\bar{\chi}_\circ v - \bar{\psi}_\circ v, \\ H_\circ \bar{\chi}_\circ \bar{\psi}_\circ v &= h\bar{\chi}_\circ \bar{\psi}_\circ v, & H_\circ \bar{\psi}_\circ v &= -(h + \frac{3k^2}{2} \Delta + \frac{1}{4} (k^2 - \frac{1}{4})) \bar{\chi}_\circ v + (h + \frac{1}{2}) \bar{\psi}_\circ v. \end{aligned}$$

The eigenvalues of H_0 on the subspace spanned by $\bar{\chi}_0 v$ and $\bar{\psi}_0 v$ are thus $h + \frac{1}{4} \pm \sqrt{h + \frac{1}{4}(6\Delta + 1)k^2}$. This action is therefore also nondiagonalisable when $h = -\frac{1}{4}(6\Delta + 1)k^2$.

This classifies irreducible $\text{Zhu}[W_{\text{pr}}^k]$ -modules with a weight vector. Could there exist an irreducible $\text{Zhu}[W_{\text{pr}}^k]$ -module without a weight vector? To tackle this question, recall that because $\text{Zhu}[W_{\text{pr}}^k]$ has countable dimension, its irreducible modules do too. Dixmier's lemma therefore shows that L_0 always acts as a multiple of the identity on any irreducible.

Choose $v \neq 0$ in an irreducible $\text{Zhu}[W_{\text{pr}}^k]$ -module. By acting with χ_0 and ψ_0 , we may assume without loss of generality that v is annihilated by both χ_0 and ψ_0 . If v is a weight vector, then our irreducible is a highest-weight module by Lemma 3.1. So suppose that v is not weight. As $H_0 v$ is then nonzero, the submodule it generates must still contain v , by irreducibility. Because we can order monomials in $\text{Zhu}[W_{\text{pr}}^k]$, this submodule is spanned by monomials of the form $(\bar{\psi}_0)^\ell (\bar{\chi}_0)^m H_0^n v$, with $n \geq 1$. However, our irreducible admits a grading inherited from that of $\text{Zhu}[W_{\text{pr}}^k]$ by setting the grade of v to 0. To obtain v , we may thus restrict to monomials with $\ell = m = 0$, concluding that $v = p(H_0)v$ for some polynomial p with zero constant term. But now, H_0 preserves the finite-dimensional subspace $\{H_0^n v : n = 0, 1, \dots, \deg(p) - 1\}$ and so this subspace contains a weight vector. We therefore conclude that every irreducible has a weight vector.

Proposition 3.3. *Every irreducible $\text{Zhu}[W_{\text{pr}}^k]$ -module is highest weight.*

Recall that a finitely generated module over a vertex-operator superalgebra is lower bounded if its conformal weights have a minimal real part. The subspace realising this minimum is called the top space, and it is naturally a module for the Zhu algebra. We define a weight vector in a W_{pr}^k -module to be an eigenvector of H_0 that is also a generalised eigenvector of L_0 , its weight then being the pair (h, Δ) of eigenvalues of H_0 and L_0 (respectively). A weight vector is a highest-weight vector if it is also annihilated by χ_0 , ψ_0 and all the modes of $\widehat{\mathfrak{psl}}_{2|2}$ with positive indices. The previous proposition now has the following consequence for the representation theory of W_{pr}^k .

Theorem 3.4. *Every irreducible lower-bounded W_{pr}^k -module is highest weight, hence determined up to isomorphism by its highest weight (h, Δ) . Moreover, the top space of the irreducible is 1-dimensional, if $\Delta = 0$, and is otherwise 4-dimensional.*

4. INVERSE REDUCTION AND THE $N = 4$ SUPERCONFORMAL ALGEBRA

We conclude by explaining how the paradigm of inverse quantum hamiltonian reduction can be used to transfer representation-theoretic information between W-algebras. For this, we explicitly construct a so-called Semikhatov realisation of the small $N = 4$ superconformal vertex-operator superalgebra W_{min}^k from W_{pr}^k and sketch how to determine when it descends to $W_{\text{k}}^{\text{min}}$ and W_{k}^{pr} . As the representation theory of W_{pr}^k and W_{k}^{pr} is still mysterious for general levels k , we then restrict to $k = \pm \frac{1}{2}$. The corresponding $N = 4$ central charges are -3 and -9 , both of which have been previously considered in the literature. In particular, inverse reduction allows us to easily recover the $c = -9$ results of [1, 5]. We take the opportunity to complement these known results with similar ones for $c = -3$ and, for both central charges, detailed structures for certain nonsemisimple $W_{\pm 1/2}^{\text{min}}$ -modules, some of which are logarithmic.

4.1. Inverse quantum hamiltonian reduction. As previously mentioned, the quantum hamiltonian reduction W_{min}^k of $V^k(\mathfrak{psl}_{2|2})$ with respect to a minimal nilpotent element is isomorphic to the small $N = 4$ superconformal vertex-operator superalgebra. In the framework of inverse quantum hamiltonian reduction [2, 48], we expect a Semikhatov realisation of W_{min}^k in terms of W_{pr}^k and some free field algebra F :

$$(4.1) \quad W_{\text{min}}^k \hookrightarrow W_{\text{pr}}^k \otimes F.$$

This embedding should moreover descend to an embedding involving the simple quotients $W_{\text{k}}^{\text{min}}$ and W_{k}^{pr} , at least for most k , and thereby provide nontrivial insights into the representation theory of W_{min}^k and $W_{\text{k}}^{\text{min}}$.

To construct such a Semikhatov realisation, we will utilise an explicit presentation of W_{min}^k , following [38, 39]. This has the following features:

- A strong generating set consists of even elements J^+ , J^0 , J^- and T , as well as odd elements G^+ , G^- , $\bar{G}^+$ and $\bar{G}^-$.
- T is a conformal vector for W_{min}^k of central charge $-6(k + 1)$ and the corresponding conformal weights of the strong generators labelled by J are 1, while those of the strong generators labelled by G are $\frac{3}{2}$.
- All the strongly generating fields except $T(z)$ are primary with respect to this conformal structure.

- The zero modes of J^+ , J^0 and J^- form an $\mathfrak{sl}_2$ -triple and the vertex subalgebra they generate is isomorphic to the affine vertex algebra $V^{-k-1}(\mathfrak{sl}_2)$.
- G^+ and G^- span a fundamental module of $\mathfrak{sl}_2$, as do $\bar{G}^+$ and $\bar{G}^-$. Here, the labels $+$ and $-$ correspond to the eigenvalue of J^0 . We normalise these $\mathfrak{sl}_2$ -actions so that $J_0^\pm G^\mp = G^\pm$ and $J_0^\pm \bar{G}^\mp = \bar{G}^\pm$.

Given this information, it only remains to specify the operator product expansions involving $G^\pm$ and $\bar{G}^\pm$, the nonregular of which are given by

$$(4.2) \quad G^\pm(z)\bar{G}^\pm(w) \sim \pm \frac{2J^\pm(w)}{(z-w)^2} \pm \frac{\partial J^\pm(w)}{z-w}, \quad G^\pm(z)\bar{G}^\mp(w) \sim \pm \frac{2(k+1)\mathbb{1}}{(z-w)^3} - \frac{J^0(w)}{(z-w)^2} - \frac{\frac{1}{2}\partial J^0(w) \pm T(w)}{z-w}.$$

We remark that if one requires that the action of the even generating fields is invariant under a 2π -rotation about the origin, while that of the odd fields need only be invariant under a 4π -rotation, then the representation theories of $W_{\min}^k$ and $W_k^{\min}$ decompose into two sectors: the Neveu–Schwarz sector in which all odd fields are invariant and the Ramond sector in which all odd fields are negated under a 2π -rotation. These two sectors are moreover equivalent as categories because there exist spectral flow maps σ^ℓ , parametrised for $\ell \in \frac{1}{2}\mathbb{Z}$ by ℓJ^0 (as in [43]). At the level of the modes of the generating fields, σ^ℓ adds $\mp\ell$ to the indices of the $G^\pm$ - and $\bar{G}^\pm$ -modes, while it adds $\mp 2\ell$ to those of the $J^\pm$ -modes. Otherwise, we have

$$(4.3) \quad \sigma^\ell(J_n^0) = J_n^0 + 2(k+1)\ell\delta_{n,0}\mathbb{1} \quad \text{and} \quad \sigma^\ell(T_n) = T_n - \ell J_n^0 - (k+1)\ell^2\delta_{n,0}\mathbb{1}.$$

There is also the conjugation automorphism w of the mode algebra of $W_{\min}^k$. It preserves the mode indices and T , while acting as the Weyl reflection of $\mathfrak{sl}_2$ on J^0 and $J^\pm$. The action on the odd generators is given by

$$(4.4) \quad w(G^\pm) = G^\mp \quad \text{and} \quad w(\bar{G}^\pm) = -\bar{G}^\mp.$$

Let $F = \Pi$ be the Friedan–Martinec–Shenker bosonisation of the bosonic ghosts vertex algebra, also known as the half-lattice vertex algebra. This has strong generators c , d and e^{mc} , for $m \in \mathbb{Z}$, whose nonregular operator product expansions are [32]

$$(4.5) \quad c(z)d(w) \cong \frac{2\mathbb{1}}{(z-w)^2}, \quad d(z)e^{mc}(w) \sim \frac{2me^{mc}(w)}{z-w}.$$

We also introduce the following convenient alternative basis for the Heisenberg elements in Π :

$$(4.6) \quad a = +\frac{1}{2}(k+1)c + \frac{1}{4}d, \quad b = -\frac{1}{2}(k+1)c + \frac{1}{4}d.$$

It is easy to check that $a(z)b(w)$ is regular. Finally, we equip Π with the conformal vector

$$(4.7) \quad t = \frac{1}{2}:cd: - \partial a.$$

It corresponds to central charge $-2(3k+2)$. The conformal weights of c , d and e^{mc} are 1, 1 and $\frac{1}{2}m$, respectively.

Given this setup, it is easy to verify (e.g. using OPEDEFS) the following result.

Theorem 4.1. *For $k \neq 0$, there is a homomorphism $\Phi^k: W_{\min}^k \rightarrow W_{\text{pr}}^k \otimes \Pi$ of vertex-operator algebras, determined by*

$$(4.8) \quad \begin{aligned} \Phi^k(J^+) &= e^{2c}, & \Phi^k(J^0) &= 2b, & \Phi^k(T) &= L + t, & \Phi^k(G^+) &= :\chi e^c:, & \Phi^k(\bar{G}^+) &= : \bar{\chi} e^c:, \\ \Phi^k(G^-) &= -:(\psi + \chi a - \frac{1}{2}(k + \frac{1}{2})\partial\chi)e^{-c}:, & \Phi^k(\bar{G}^-) &= -:(\bar{\psi} + \bar{\chi} a - \frac{1}{2}(k + \frac{1}{2})\partial\bar{\chi})e^{-c}:, \\ \Phi^k(J^-) &= :(H - \frac{1}{2}(k-1)S - aa - \frac{1}{2}(k + \frac{1}{2})(3k-1)\chi\bar{\chi} + k\partial a)e^{-2c}:. \end{aligned}$$

To convert this into a Semikhatov realisation of $W_{\min}^k$, we need to demonstrate that Φ^k is injective. This is clear for generic levels, but is rather subtle otherwise. However, it follows from a combinatorial argument based on the “triangularity” of Φ^k , see [7] for an example of such an argument. Here, triangularity refers to the fact that the strong generators of $W_{\min}^k$ and $W_{\text{pr}}^k \otimes \Pi$ may be ordered so that the image of the n -th generator of $W_{\min}^k$ is expressed in terms of the m -th generators, with $m \leq n$. (For Π , we only need to consider d and e^c for this purpose, see [7].) We omit the details.

It remains to obtain a Semikhatov realisation of $W_k^{\min}$. As in [7], this follows from a remarkable property of inverse quantum hamiltonian reduction. First, recall [13] that Π admits a family of irreducible lower-bounded modules

$$(4.9) \quad \Pi_{[\lambda]} = \Pi e^{-a+\lambda c}, \quad [\lambda] \in \mathbb{C}/\mathbb{Z}.$$

The top space of $\Pi_{[\lambda]}$ is spanned by the $e^{-a+\mu c}$ with $\mu \in [\lambda]$ and it has conformal weight $-\frac{1}{4}(k+1)$. Note that because e^c has conformal weight $\frac{1}{2}$, the $\Pi_{[\lambda]}$ are actually $\mathbb{Z}_2$ -twisted Π -modules.

With these (twisted) Π -modules, we introduce the Adamović functors

$$(4.10) \quad \text{Ad}_{[\lambda]} = \text{Res}_{W_{\min}^k}^{W_{\text{pr}}^k \otimes \Pi} (- \otimes \Pi_{[\lambda]}), \quad [\lambda] \in \mathbb{C}/\mathbb{Z}.$$

The remarkable property is now that if $\mathcal{M}$ is a simple W_{pr}^k -module, then $\text{Ad}_{[\lambda]}(\mathcal{M})$ is almost irreducible. This means that it is lower bounded, generated by its top space, and is such that every nonzero submodule has nonzero intersection with the top space. We will also omit the details that establish this property, referring again to [7] for a similar example.

This property of Adamović functors has many important applications. One such is the precise condition under which Φ^k descends to a Semikhatov realisation

$$(4.11) \quad \Phi_k : W_k^{\min} \hookrightarrow W_k^{\text{pr}} \otimes \Pi$$

of the simple quotients. The idea is as follows [7]. Since W_k^{pr} is irreducible, $\text{Ad}_{[\lambda]}(W_k^{\text{pr}})$ is an almost-irreducible $W_{\min}^k$ -module, hence any of its nonzero submodules intersect its top space nontrivially. But, for a suitable choice of $[\lambda]$, this module is a spectral flow of $W_k^{\text{pr}} \otimes \Pi$. It follows that the image of the maximal submodule of $W_{\min}^k$ is either 0 or it has nonzero intersection with the “spectrally flown top space” $\text{span}\{(J^+)^n|0\rangle : n \in \mathbb{Z}\}$ of $W_k^{\text{pr}} \otimes \Pi$. As $J^+ \in V^{-k-1}(\mathfrak{sl}_2)$, the latter is impossible unless $k \in \mathbb{Z}_{\leq -1}$.

Theorem 4.2. *For $k \notin \mathbb{Z}_{\leq 0}$, the embedding of (4.8) induces the simple Semikhatov realisation (4.11).*

4.2. Constructing weight $W_{\pm 1/2}^{\min}$ -modules. Recall that the simple principal W-algebra W_k^{pr} collapses to the symplectic fermions vertex-operator superalgebra SF when the level is $k = \pm \frac{1}{2}$. The corresponding simple minimal W-algebras $W_{1/2}^{\min}$ and $W_{-1/2}^{\min}$ have central charges -9 and -3 , respectively. For convenience, we specialise the Semikhatov realisation of Theorems 4.1 and 4.2 to these levels by setting H, S, ψ and $\bar{\psi}$ to 0.

Corollary 4.3. *For $k = \pm \frac{1}{2}$, there is a Semikhatov realisation $\Phi_k : W_k^{\min} \hookrightarrow \text{SF} \otimes \Pi$ given by*

$$(4.12) \quad \begin{aligned} \Phi_k(J^+) &= e^{2c}, & \Phi_k(J^0) &= 2b, & \Phi_k(J^-) &= -(aa - k\partial a)e^{-2c} - \frac{1}{2}(k + \frac{1}{2})(3k - 1) : \chi \bar{\chi} : e^{-2c}, \\ \Phi_k(G^+) &= \chi e^c, & \Phi_k(G^-) &= \frac{1}{2}(k + \frac{1}{2})\partial \chi e^{-c} - \chi : a e^{-c} :, & \Phi_k(T) &= \frac{1}{2} : cd : - \partial a - \frac{1}{2} : \chi \bar{\chi} :, \\ \Phi_k(\bar{G}^+) &= \bar{\chi} e^c, & \Phi_k(\bar{G}^-) &= \frac{1}{2}(k + \frac{1}{2})\partial \bar{\chi} e^{-c} - \bar{\chi} : a e^{-c} :, \end{aligned}$$

These free-field realisations of the $c = -9$ and $c = -3$ (small) $N = 4$ superconformal algebra are closely related to other realisations that have appeared in the literature.

- For $W_{1/2}^{\min}$ ($c = -9$), [1, Prop. 5.1 & Thm. 6.1] replaces SF and Π with the $c = -11$ fermionic ghost vertex-operator superalgebra FG and the $c = 2$ bosonic ghost vertex-operator algebra BG, respectively. We remark that $\text{SF} \subset \text{FG}$ and $\Pi \supset \text{BG}$ as vertex superalgebras (these are not conformal embeddings).
- In [5, Prop. 1], $W_{1/2}^{\min}$ ($c = -9$) is shown to embed into a $\mathbb{Z}_2$ -orbifold of $\text{SF} \otimes \Pi^{1/2}$, where $\Pi^{1/2}$ is a simple-current extension of Π . This is very close to Corollary 4.3, but is very different in flavour: their realisation uses screening operators and does not identify SF as a W-algebra.
- On the other hand, [16, Thm. 4.14] realises $W_{-1/2}^{\min}$ ($c = -3$) as a simple-current extension of the tensor product of the $c = -2$ triplet vertex-operator algebra $W(2)$ and the $c = -1$ affine vertex-operator algebra $L_{-1/2}(\mathfrak{sl}_2)$. This is also close to Corollary 4.3: $W(2) \subset \text{SF}$ and $L_{-1/2}(\mathfrak{sl}_2) \subset \text{BG} \subset \Pi$ as vertex-operator superalgebras (these are conformal embeddings).

Recall that SF has a single irreducible untwisted (hence Neveu–Schwarz) module $\mathcal{S}^{\text{NS}} \cong \text{SF}$ and a single irreducible $\mathbb{Z}_2$ -twisted (hence Ramond) module $\mathcal{S}^{\text{R}}$. The highest-weight vectors of $\mathcal{S}^{\text{NS}}$ and $\mathcal{S}^{\text{R}}$, which we denote by $|\text{NS}\rangle$ and $|\text{R}\rangle$, have conformal weights 0 and $-\frac{1}{8}$, respectively. The Adamović functors thus give two families of almost-irreducible (twisted) $W_{\pm 1/2}^{\min}$ -modules, parametrised by $[\lambda] \in \mathbb{C}/\mathbb{Z}$:

$$(4.13) \quad \mathcal{N}_{[\lambda]}^{\text{R}} = \text{Ad}_{[\lambda]}(\mathcal{S}^{\text{NS}}) \quad \text{and} \quad \mathcal{N}_{[\lambda]}^{\text{NS}} = \text{Ad}_{[\lambda]}(\mathcal{S}^{\text{R}}).$$

Note that because $\Pi_{[\lambda]}$ is $\mathbb{Z}_2$ -twisted, the $\mathcal{N}_{[\lambda]}^R$ are too — they belong to the Ramond sector of $W_{\pm 1/2}^{\min}$. The $\mathcal{N}_{[\lambda]}^{\text{NS}}$ are untwisted, hence Neveu–Schwarz.

Recall the characters and supercharacters of the irreducible SF-modules:

$$\begin{aligned}
 \text{ch}^-[\mathcal{S}^{\text{NS}}](y; q) &= q^{1/12} \prod_{n=1}^{\infty} (1 - yq^n)(1 - y^{-1}q^n) = \frac{1}{y^{1/2} - y^{-1/2}} \frac{i\vartheta_1(y; q)}{\eta(q)}, \\
 \text{ch}^+[\mathcal{S}^{\text{NS}}](y; q) &= q^{1/12} \prod_{n=1}^{\infty} (1 + yq^n)(1 + y^{-1}q^n) = \frac{1}{y^{1/2} + y^{-1/2}} \frac{\vartheta_2(y; q)}{\eta(q)}, \\
 \text{ch}^+[\mathcal{S}^R](y; q) &= q^{-1/24} \prod_{n=1}^{\infty} (1 + yq^{n-1/2})(1 + y^{-1}q^{n-1/2}) = \frac{\vartheta_3(y; q)}{\eta(q)}, \\
 \text{ch}^-[\mathcal{S}^R](y; q) &= q^{-1/24} \prod_{n=1}^{\infty} (1 - yq^{n-1/2})(1 - y^{-1}q^{n-1/2}) = \frac{\vartheta_4(y; q)}{\eta(q)}.
 \end{aligned}
 \tag{4.14}$$

Here, we have included the horizontal grading as in (2.27). For the lower-bounded Π -modules $\Pi_{[\lambda]}$, we define characters as follows:

$$\text{ch}^+[\Pi_{[\lambda]}](z; q) = \text{tr}_{\Pi_{[\lambda]}} z^{2b_0} q^{t_0 + (3k+2)/12} = \frac{q^{-(k+1)/4 + (3k+2)/12}}{\prod_{n=1}^{\infty} (1 - q^n)^2} \sum_{n \in \mathbb{Z}} z^{\lambda+n} = z^{\lambda} \frac{\delta(z)}{\eta(q)^2}.
 \tag{4.15}$$

Here, $\delta(z)$ is the formal series (or distribution) $\sum_{n \in \mathbb{Z}} z^n$.

Because Corollary 4.3 identifies J^0 and $2b$, we may define graded (super)characters of $W_{\pm 1/2}^{\min}$ -modules that keep track of the horizontal grading (y), the J_0^0 -eigenvalue (z) and the conformal weight (q). For the images (4.13) under the Adamović functors, these (super)characters have the simple forms

$$\begin{aligned}
 \text{ch}^-[\mathcal{N}_{[\lambda]}^R](y, z; q) &= \frac{z^{\lambda}}{y^{1/2} - y^{-1/2}} \frac{i\vartheta_1(y; q)\delta(z)}{\eta(q)^3}, & \text{ch}^+[\mathcal{N}_{[\lambda]}^{\text{NS}}](y, z; q) &= z^{\lambda} \frac{\vartheta_3(y; q)\delta(z)}{\eta(q)^3}, \\
 \text{ch}^+[\mathcal{N}_{[\lambda]}^R](y, z; q) &= \frac{z^{\lambda}}{y^{1/2} + y^{-1/2}} \frac{\vartheta_2(y; q)\delta(z)}{\eta(q)^3}, & \text{ch}^-[\mathcal{N}_{[\lambda]}^{\text{NS}}](y, z; q) &= z^{\lambda} \frac{\vartheta_4(y; q)\delta(z)}{\eta(q)^3}.
 \end{aligned}
 \tag{4.16}$$

Given these results, we may also use (4.3) to determine the (super)characters of the spectral flows of these modules.

4.3. Degenerations and spectral flows. Consider now the almost-irreducible $W_{\pm 1/2}^{\min}$ -modules $\mathcal{N}_{[\lambda]}^R$. These belong to the Ramond sector and so are completely determined by the action of the zero modes, including $G_0^{\pm}$ and $\bar{G}_0^{\pm}$, on their top spaces [20]. A basis for this top space is given by the vectors

$$|\mu\rangle^R = |\text{NS}\rangle \otimes e^{-a+\mu c}, \quad \mu \in [\lambda].
 \tag{4.17}$$

Using Corollary 4.3, we compute the action of the generators' zero modes on these top space vectors. The even modes' actions are given by

$$\begin{aligned}
 J_0^+ |\mu\rangle^R &= |\mu + 2\rangle^R, & J_0^0 |\mu\rangle^R &= \mu |\mu\rangle^R, & T_0 |\mu\rangle^R &= -\frac{1}{4}(k+1) |\mu\rangle^R, \\
 J_0^- |\mu\rangle^R &= -\frac{1}{4}(\mu - k - 1)(\mu + k - 1) |\mu - 2\rangle^R,
 \end{aligned}
 \tag{4.18}$$

while the odd modes all act as 0. The top space of $\mathcal{N}_{[\lambda]}^R$ thus decomposes as a direct sum of two (generically irreducible) modules for the zero-mode algebra (the Ramond-twisted Zhu algebra) of $W_{\pm 1/2}^{\min}$. We may therefore write

$$\mathcal{N}_{[\lambda]}^R = \mathcal{M}_{[\lambda]}^R \oplus \mathcal{M}_{[\lambda+1]}^R, \quad [\lambda] = \lambda + \mathbb{Z} \in \mathbb{C}/\mathbb{Z},
 \tag{4.19}$$

where $[\lambda] = \lambda + 2\mathbb{Z} \in \mathbb{C}/2\mathbb{Z}$ and $\mathcal{M}_{[\lambda]}^R$ denotes the submodule of $\mathcal{N}_{[\lambda]}^R$ generated by the $|\mu\rangle^R$ with $\mu \in [\lambda]$.

Note that the top spaces of the $\mathcal{M}_{[\lambda]}^R$, $[\lambda] \in \mathbb{C}/2\mathbb{Z}$, may be viewed as dense modules for the even zero-mode subalgebra of $W_{\pm 1/2}^{\min}$ (which is isomorphic to $\mathfrak{gl}_2$). Moreover, the (super)character of $\mathcal{M}_{[\lambda]}^R$ is obtained from that of $\mathcal{N}_{[\lambda]}^R$, given in (4.16), by replacing $\delta(z)$ by $\delta(z^2)$.

It follows immediately from (4.18) that $\mathcal{M}_{[\lambda]}^R$ is reducible if and only if there exists a conjugate highest-weight vector $|\mu\rangle^R$ with $\mu \in [\lambda]$. Moreover, such a vector only exists when $\mu = 1 \pm k = \frac{1}{2}, \frac{3}{2}$. We thus say that the family $\{\mathcal{M}_{[\lambda]}^R : [\lambda] \in \mathbb{C}/2\mathbb{Z}\}$ degenerates at $[\lambda] = [\pm \frac{1}{2}]$.

Let $\mathcal{L}_{\mu}^R$ denote the irreducible highest-weight $W_{\pm 1/2}^{\min}$ -module, in the Ramond sector, whose highest-weight vector has J_0^0 -eigenvalue μ and T_0 -eigenvalue $-\frac{1}{4}(k+1)$. Recalling the conjugation automorphism w from Section 4.1, the reducible

cases are then characterised by the following nonsplit short exact sequences:

$$(4.20) \quad 0 \longrightarrow w(\mathcal{L}_{-1/2}^R) \longrightarrow \mathcal{M}_{\llbracket 1/2 \rrbracket}^R \longrightarrow \mathcal{L}_{-3/2}^R \longrightarrow 0, \quad 0 \longrightarrow w(\mathcal{L}_{-3/2}^R) \longrightarrow \mathcal{M}_{\llbracket -1/2 \rrbracket}^R \longrightarrow \mathcal{L}_{-1/2}^R \longrightarrow 0.$$

This result deserves a few comments.

- The $\mathcal{M}_{\llbracket \lambda \rrbracket}^R$ are almost irreducible, see for example [27, Prop. 4.8].
- The submodules in (4.20) are irreducible because any nonzero proper submodule would also be one of $\mathcal{M}_{\llbracket \pm 1/2 \rrbracket}^R$, hence would intersect the top space.
- The quotients are not obviously irreducible. This follows easily if one has a classification of highest-weight modules, such as [1, Prop. 6.2] for $k = \frac{1}{2}$. Otherwise, the irreducibility may be established using the methods developed in [42, Prop. 4.13], which also demonstrate exactness.
- To our knowledge, this is the first time the images of the Adamović functors have produced modules that are generically completely reducible rather than generically irreducible. In this case, this follows from an observation made in [5], whereby $W_{\pm 1/2}^{\min}$ in fact embeds into the even subalgebra of $SF \otimes \Pi$ with respect to a $\mathbb{Z}_2 \times \mathbb{Z}_2$ action. The $\mathbb{Z}_2$ action on SF is the usual action on even and odd subspaces. The action on Π is on the lattice generators e^{mc} and is determined by whether m is even or odd. It is clear from inspecting (4.3) that even generators of $W_{\pm 1/2}^{\min}$ are mapped into the $(0, 0)$ subspace, and odd elements of $W_{\pm 1/2}^{\min}$ are mapped into the $(1, 1)$ subspace with respect to this action.

One can repeat the above analysis in the Neveu–Schwarz sector. Now, the top space of $\mathcal{N}_{\llbracket \lambda \rrbracket}^{\text{NS}}$, $[\lambda] \in \mathbb{C}/\mathbb{Z}$, has the following basis vectors:

$$(4.21) \quad |\mu\rangle^{\text{NS}} = |\mathbf{R}\rangle \otimes e^{-a+\mu c}, \quad \mu \in [\lambda].$$

The odd generators have no zero modes, and the even zero modes' actions are given explicitly by

$$(4.22) \quad \begin{aligned} J_0^+ |\mu\rangle^{\text{NS}} &= |\mu + 2\rangle^{\text{NS}}, & J_0^0 |\mu\rangle^{\text{NS}} &= \mu |\mu\rangle^{\text{NS}}, & T_0 |\mu\rangle^{\text{NS}} &= -\frac{1}{4}(k + \frac{3}{2}) |\mu\rangle^{\text{NS}}, \\ J_0^- |\mu\rangle^{\text{NS}} &= -(\frac{1}{4}(\mu - k - 1)(\mu + k - 1) + \frac{1}{8}(k + \frac{1}{2})(3k - 1)) |\mu - 2\rangle^{\text{NS}}, \end{aligned} \quad \mu \in [\lambda].$$

We therefore again get a direct sum decomposition:

$$(4.23) \quad \mathcal{N}_{\llbracket \lambda \rrbracket}^{\text{NS}} = \mathcal{M}_{\llbracket \lambda \rrbracket}^{\text{NS}} \oplus \mathcal{M}_{\llbracket \lambda + 1 \rrbracket}^{\text{NS}}.$$

Let $\mathcal{M}_{\llbracket \mu \rrbracket}^{\text{NS}}$ denote the irreducible highest-weight $W_{\pm 1/2}^{\min}$ -module, in the Neveu–Schwarz sector, whose highest-weight vector has J_0^0 -eigenvalue μ and T_0 -eigenvalue $-\frac{1}{4}(k + \frac{3}{2})$.

When $k = -\frac{1}{2}$, the Neveu–Schwarz analysis is almost identical to the Ramond case. The family $\{\mathcal{M}_{\llbracket \lambda \rrbracket}^{\text{NS}} : \llbracket \lambda \rrbracket \in \mathbb{C}/2\mathbb{Z}\}$ degenerates at $\llbracket \lambda \rrbracket = \llbracket \pm \frac{1}{2} \rrbracket$ and there we have the following nonsplit short exact sequences:

$$(4.24) \quad 0 \longrightarrow w(\mathcal{L}_{-1/2}^{\text{NS}}) \longrightarrow \mathcal{M}_{\llbracket 1/2 \rrbracket}^{\text{NS}} \longrightarrow \mathcal{L}_{-3/2}^{\text{NS}} \longrightarrow 0, \quad 0 \longrightarrow w(\mathcal{L}_{-3/2}^{\text{NS}}) \longrightarrow \mathcal{M}_{\llbracket -1/2 \rrbracket}^{\text{NS}} \longrightarrow \mathcal{L}_{-1/2}^{\text{NS}} \longrightarrow 0.$$

The analysis for $k = \frac{1}{2}$ is significantly more interesting. In this case, $J_0^- |\mu\rangle^{\text{NS}} = 0$ has only one solution: $\mu = 1$. The family $\{\mathcal{M}_{\llbracket \lambda \rrbracket}^{\text{NS}} : \llbracket \lambda \rrbracket \in \mathbb{C}/2\mathbb{Z}\}$ thus only degenerates at $\llbracket \lambda \rrbracket = \llbracket 1 \rrbracket$. But, $\mathcal{M}_{\llbracket 1 \rrbracket}^{\text{NS}}$ has more than two composition factors! Indeed, its Loewy diagram is

$$(4.25) \quad \begin{array}{ccc} & \mathcal{L}_{-1}^{\text{NS}} & \\ & \swarrow \quad \searrow & \\ \mathcal{L}_0^{\text{NS}} & \mathcal{M}_{\llbracket 1 \rrbracket}^{\text{NS}} & \mathcal{L}_0^{\text{NS}} \\ & \swarrow \quad \searrow & \\ & w(\mathcal{L}_{-1}^{\text{NS}}) & \end{array}$$

This structure is also indicated schematically in Figure 2.

We make a few further comments on the structure of $\mathcal{M}_{\llbracket 1 \rrbracket}^{\text{NS}}$.

- This module is not logarithmic: T_0 acts semisimply (its head and socle are not isomorphic).
- It is well known that an almost-irreducible module may have composition factors that are not detected by the top space, see [42, §4.5]. For simple affine vertex algebras (and W-algebras), this appears to be a feature of the representation




FIGURE 2. An illustration of the structure of the $W_{1/2}^{\min}$ -module $\mathcal{M}_{[1]}^{\text{NS}}$. The dots represent weight vectors, with the J_0^0 -eigenvalue increasing from left to right and the T_0 -eigenvalue Δ from top to bottom.

theory at nonadmissible levels. Given that the level of the affine vertex subalgebra generated by J^0 and $J^\pm$ is admissible for $k = -\frac{1}{2}$, but nonadmissible for $k = \frac{1}{2}$, this qualitative structural difference should perhaps not be surprising.

- It is nevertheless quite satisfying to use (4.12) to explicitly verify that $G_{-1/2}^-$ and $\bar{G}_{-1/2}^-$ annihilate $|1\rangle^{\text{NS}}$, while $G_{1/2}^-$ and $\bar{G}_{1/2}^-$ annihilate both $G_{-1/2}^+|-1\rangle^{\text{NS}}$ and $\bar{G}_{-1/2}^+|-1\rangle^{\text{NS}}$.
- The fact that $\mathcal{M}_{[1]}^{\text{NS}}$ does not have more than four composition factors uses the classification [1, Prop. 6.2] of irreducible highest-weight $W_{1/2}^{\min}$ -modules (there are only two: $\mathcal{L}_0^{\text{NS}}$ and $\mathcal{L}_{-1}^{\text{NS}}$). The inverse-reduction technology developed in [8] can also be used to prove this classification result, though we shall not do so here.
- As noted above, $W_{1/2}^{\min}$ is a vertex subalgebra of the tensor product of a $c = -11$ fermionic ghost system and a $c = 2$ bosonic ghost system. In our language, [1, Prop. 6.3] identifies this tensor product as a nonsplit extension of $\mathcal{L}_{-1}^{\text{NS}}$ by $\mathcal{L}_0^{\text{NS}}$, hence as a length-2 quotient of $\mathcal{M}_{[1]}^{\text{NS}}$.
- It is worth remarking that the quotient $\mathcal{M}_{[1]}^{\text{NS}}/w(\mathcal{L}_{-1}^{\text{NS}})$ is a highest-weight $W_{1/2}^{\min}$ -module with two linearly independent singular vectors that share the same J_0^0 - and T^0 -eigenvalues.

We conclude by reporting the action of spectral flow on the irreducible highest-weight $W_{\pm 1/2}^{\min}$ -modules. For $k = -\frac{1}{2}$, the spectral flow orbits feature ordinary modules that do not appear as composition factors of the almost-irreducible modules constructed by inverse reduction. These follow from explicit computation using (4.3):

$$(4.26) \quad \begin{aligned} \dots &\xrightarrow{\sigma^{1/2}} w(\mathcal{L}_{-1/2}^{\text{R}}) \xrightarrow{\sigma^{1/2}} \mathcal{L}_0^{\text{NS}} \xrightarrow{\sigma^{1/2}} \mathcal{L}_{-1/2}^{\text{R}} \xrightarrow{\sigma^{1/2}} \dots, & \dots &\xrightarrow{\sigma^{1/2}} w(\mathcal{L}_{-1/2}^{\text{NS}}) \xrightarrow{\sigma^{1/2}} \mathcal{L}_0^{\text{R}} \xrightarrow{\sigma^{1/2}} \mathcal{L}_{-1/2}^{\text{NS}} \xrightarrow{\sigma^{1/2}} \dots, \\ \dots &\xrightarrow{\sigma^{1/2}} w(\mathcal{L}_{-3/2}^{\text{R}}) \xrightarrow{\sigma^{1/2}} \mathcal{L}_1^{\text{NS}} \xrightarrow{\sigma^{1/2}} \mathcal{L}_{-3/2}^{\text{R}} \xrightarrow{\sigma^{1/2}} \dots, & \dots &\xrightarrow{\sigma^{1/2}} w(\mathcal{L}_{-3/2}^{\text{NS}}) \xrightarrow{\sigma^{1/2}} \mathcal{L}_1^{\text{R}} \xrightarrow{\sigma^{1/2}} \mathcal{L}_{-3/2}^{\text{NS}} \xrightarrow{\sigma^{1/2}} \dots. \end{aligned}$$

Here, the dots indicate irreducible modules that are not lower bounded. We also note the similarity to the spectral flow orbits of $L_{-1/2}(\mathfrak{sl}_2)$, see [46, Fig. 3]. Contrarily, for $k = \frac{1}{2}$, the orbits contain no new lower-bounded modules:

$$(4.27) \quad \begin{aligned} \dots &\xrightarrow{\sigma^{1/2}} w(\mathcal{L}_{-3/2}^{\text{R}}) \xrightarrow{\sigma^{1/2}} \mathcal{L}_0^{\text{NS}} \xrightarrow{\sigma^{1/2}} \mathcal{L}_{-3/2}^{\text{R}} \xrightarrow{\sigma^{1/2}} \dots, \\ \dots &\xrightarrow{\sigma^{1/2}} w(\mathcal{L}_{-1/2}^{\text{R}}) \xrightarrow{\sigma^{1/2}} \mathcal{L}_{-1}^{\text{NS}} \xrightarrow{\sigma^{1/2}} \dots, & \dots &\xrightarrow{\sigma^{1/2}} w(\mathcal{L}_{-1}^{\text{NS}}) \xrightarrow{\sigma^{1/2}} \mathcal{L}_{-1/2}^{\text{R}} \xrightarrow{\sigma^{1/2}} \dots. \end{aligned}$$

4.4. Logarithmic $W_{\pm 1/2}^{\min}$ -modules. Recall that the Ramond sector of the symplectic fermions vertex-operator superalgebra SF is semisimple, but the Neveu–Schwarz sector is not. In particular, it contains a logarithmic module $\mathcal{S}_{(4)}^{\text{NS}}$. This is induced from the projective module of $\mathfrak{psl}_{2|2}$ and is thus an indecomposable sum of four copies of the vacuum module. We present its Loewy diagram and a basis for its top space:

$$(4.28) \quad \begin{array}{c} \mathcal{S}^{\text{NS}} \\ \swarrow \quad \searrow \\ \mathcal{S}^{\text{NS}} \quad \mathcal{S}_{(4)}^{\text{NS}} \quad \mathcal{S}^{\text{NS}} \\ \searrow \quad \swarrow \\ \mathcal{S}^{\text{NS}} \end{array} \quad \begin{array}{c} |\text{NS}; t\rangle \\ \swarrow \quad \searrow \\ |\text{NS}; m\rangle \quad |\text{NS}; \bar{m}\rangle \\ \swarrow \quad \searrow \\ |\text{NS}; b\rangle \end{array} \quad \begin{array}{c} \chi_0 \\ \bar{\chi}_0 \\ -\chi_0 \end{array}$$

The logarithmic nature of $\mathcal{S}_{(4)}^{\text{NS}}$ is evident from the action of the Virasoro zero mode: it maps $|\text{NS}; t\rangle$ to $\frac{1}{2}|\text{NS}; b\rangle$.

We can apply Adamović functors to such nonsemisimple SF-modules. To warm up, we will first consider a structurally simpler example: the length-2 submodule of $\mathcal{S}_{(4)}^{\text{NS}}$ generated by $|\text{NS}; m\rangle$. This SF-module, which we shall denote by $\mathcal{S}_{(2)}^{\text{NS}}$,




FIGURE 3. An illustration of the structure of the top spaces of the $W_{\pm 1/2}^{\min}$ -modules $\mathcal{V}_{\llbracket 1/2 \rrbracket}^R$ (left) and $\mathcal{V}_{\llbracket -1/2 \rrbracket}^R$ (right). The dots represent weight vectors, with the J_0^0 -eigenvalue increasing from left to right. The top row corresponds to vectors with label “ m ” and the bottom row has label “ b ”. The shading suggests the composition factors.

has top space $\text{span}\{|\text{NS}; m\rangle, |\text{NS}; b\rangle\}$. It is moreover a self-extension of the vacuum module that coincides with a Verma module of $\widehat{\text{psl}}_{2|2}$. We shall denote its images under the Adamović functors by $\mathcal{W}_{[\lambda]}^R$, $[\lambda] \in \mathbb{C}/\mathbb{Z}$. These are lower-bounded Ramond-twisted $W_{\pm 1/2}^{\min}$ -modules.

The top space of $\mathcal{W}_{[\lambda]}^R$ has basis $\{|\mu; m\rangle^R, |\mu; b\rangle^R : \mu \in [\lambda]\}$, where we follow (4.17) but add the label “ m ” or “ b ” appropriately. Using Corollary 4.3, it is easy to determine the action of the zero-mode algebra of $W_{\pm 1/2}^{\min}$ on this basis. Indeed, the even zero modes act as in (4.18) (preserving labels), and almost all of the odd zero modes act as zero. The exceptions are those involving $\bar{\chi}_0$:

$$(4.29) \quad \bar{G}_0^+ |\mu; m\rangle^R = |\mu + 1; b\rangle^R \quad \text{and} \quad \bar{G}_0^- |\mu; m\rangle^R = -\frac{1}{2}(\mu - \frac{1}{2})|\mu - 1; b\rangle^R.$$

Because of this, we still have a direct sum decomposition

$$(4.30) \quad \mathcal{W}_{[\lambda]}^R = \mathcal{V}_{\llbracket \lambda \rrbracket}^R \oplus \mathcal{V}_{\llbracket \lambda+1 \rrbracket}^R, \quad [\lambda] \in \mathbb{C}/\mathbb{Z},$$

where $\mathcal{V}_{\llbracket \lambda \rrbracket}^R$ is spanned by the $|\mu; m\rangle^R$ and $|\mu+1; b\rangle^R$ with $\mu \in \llbracket \lambda \rrbracket$. However, the $\mathcal{V}_{\llbracket \lambda \rrbracket}^R$ are nonsemisimple and (generically) have two composition factors:

$$(4.31) \quad 0 \longrightarrow \mathcal{M}_{\llbracket \lambda+1 \rrbracket}^R \longrightarrow \mathcal{V}_{\llbracket \lambda \rrbracket}^R \longrightarrow \mathcal{M}_{\llbracket \lambda \rrbracket}^R \longrightarrow 0, \quad \llbracket \lambda \rrbracket \neq \llbracket \pm \frac{1}{2} \rrbracket.$$

When $\llbracket \lambda \rrbracket = \llbracket \pm \frac{1}{2} \rrbracket$, there is a further degeneration corresponding to the fact that $\mathcal{M}_{\llbracket \pm 1/2 \rrbracket}^R$ is reducible. In this case, J_0^- annihilates $|\mu; \bullet\rangle^R$, for $\mu = \frac{1}{2}, \frac{3}{2}$ and $\bullet = m, b$, and $\bar{G}_0^-$ annihilates $|\frac{1}{2}; m\rangle^R$. The resulting Loewy diagrams are then

$$(4.32) \quad \begin{array}{ccc} & \mathcal{L}_{-3/2}^R & \\ & \swarrow \quad \searrow & \\ \mathcal{L}_{-1/2}^R & \mathcal{V}_{\llbracket 1/2 \rrbracket}^R & w(\mathcal{L}_{-1/2}^R) \\ & \swarrow \quad \searrow & \\ & w(\mathcal{L}_{-3/2}^R) & \end{array} \quad \begin{array}{ccc} & \mathcal{L}_{-1/2}^R & \\ & \swarrow \quad \searrow & \\ \mathcal{L}_{-3/2}^R & \mathcal{V}_{\llbracket -1/2 \rrbracket}^R & w(\mathcal{L}_{-3/2}^R) \\ & \swarrow \quad \searrow & \\ & w(\mathcal{L}_{-1/2}^R) & \end{array},$$

see also Figure 3. Of course, neither module is logarithmic.

With this analysis complete, we turn to the image of the logarithmic module $\mathcal{S}_{(4)}^{\text{NS}}$ under the Adamović functor, which we denote by $\mathcal{Q}_{[\lambda]}^R$, $[\lambda] \in \mathbb{C}/\mathbb{Z}$. The top space has basis $\{|\mu; t\rangle^R, |\mu; m\rangle^R, |\mu; \bar{m}\rangle^R, |\mu; b\rangle^R : \mu \in [\lambda]\}$, where again the second label follows (4.28). On this top space, the even zero modes J_0^+ and J_0^0 act as they did on $\mathcal{W}_{[\lambda]}^R$ (preserving labels). The remaining even zero modes act, for $\bullet = t, m, \bar{m}, b$, as follows:

$$(4.33) \quad \begin{aligned} J_0^- |\mu; \bullet\rangle^R &= -\frac{1}{4}(\mu - k - 1)(\mu + k - 1)|\mu - 2; \bullet\rangle^R + \frac{1}{2}(k + \frac{1}{2})(3k - 1)|\mu - 2; b\rangle^R \delta_{\bullet, t}, \\ T_0 |\mu; \bullet\rangle^R &= -\frac{1}{4}(k + 1)|\mu; \bullet\rangle^R + \frac{1}{2}|\mu; b\rangle^R \delta_{\bullet, t}. \end{aligned}$$

The action of the odd zero modes on the top space basis vectors is again zero, except for

$$(4.34) \quad \begin{aligned} G_0^+ |\mu; t\rangle^R &= |\mu + 1; m\rangle^R, & G_0^+ |\mu; \bar{m}\rangle^R &= -|\mu + 1; b\rangle^R, \\ \bar{G}_0^+ |\mu; t\rangle^R &= |\mu + 1; \bar{m}\rangle^R, & \bar{G}_0^+ |\mu; m\rangle^R &= +|\mu + 1; b\rangle^R, \\ G_0^- |\mu; t\rangle^R &= -\frac{1}{2}(\mu - \frac{1}{2})|\mu - 1; m\rangle^R, & G_0^- |\mu; \bar{m}\rangle^R &= +\frac{1}{2}(\mu - \frac{1}{2})|\mu - 1; b\rangle^R, \\ \bar{G}_0^- |\mu; t\rangle^R &= -\frac{1}{2}(\mu - \frac{1}{2})|\mu - 1; \bar{m}\rangle^R, & \bar{G}_0^- |\mu; m\rangle^R &= -\frac{1}{2}(\mu - \frac{1}{2})|\mu - 1; b\rangle^R. \end{aligned}$$

It is clear that we once again have a direct sum decomposition

$$(4.35) \quad \mathcal{Q}_{[\lambda]}^R = \mathcal{P}_{[\lambda]}^R \oplus \mathcal{P}_{[\lambda+1]}^R, \quad [\lambda] \in \mathbb{C}/\mathbb{Z},$$

where $\mathcal{P}_{[\lambda]}^R$ is spanned by the $|\mu; t\rangle^R$, $|\mu + 1; m\rangle^R$, $|\mu + 1; \bar{m}\rangle^R$ and $|\mu; b\rangle^R$, with $\mu \in [\lambda]$. Each of these submodules is logarithmic and their Loewy diagrams have the generic form

$$(4.36) \quad \begin{array}{ccccc} & & \mathcal{M}_{[\lambda]}^R & & \\ & \swarrow & & \searrow & \\ \mathcal{M}_{[\lambda+1]}^R & & \mathcal{P}_{[\lambda]}^R & & \mathcal{M}_{[\lambda+1]}^R \\ & \swarrow & & \searrow & \\ & & \mathcal{M}_{[\lambda]}^R & & \end{array}, \quad [\lambda] \neq [\pm \frac{1}{2}].$$

It is clear from (4.33) that these modules are logarithmic.

Consider the degeneration that occurs when $[\lambda] = [\pm \frac{1}{2}]$. We will suppose first that $k = -\frac{1}{2}$. Then, the qualitative difference from the generic action given above is that J_0^- acts as zero, when $\mu = \frac{1}{2}$ or $\frac{3}{2}$, while G_0^- and $\bar{G}_0^-$ act as zero, when $\mu = \frac{1}{2}$. The resulting structures are depicted, along with the corresponding Loewy diagrams, in Figure 4.

For $k = \frac{1}{2}$, the only difference is that J_0^- no longer acts as zero when $\mu = \frac{1}{2}$ or $\frac{3}{2}$. Instead, we have

$$(4.37) \quad J_0^- |\mu; \bullet\rangle^R = \frac{1}{4} |\mu - 2; b\rangle^R \delta_{\bullet, t}.$$

This modifies the structural depictions and Loewy diagrams of Figure 4 by adding one arrow to each. Specifically, we should add an arrow in each depiction from the top dot labelled by $\frac{1}{2} (\frac{3}{2})$ to the bottom dot labelled by $-\frac{3}{2} (-\frac{1}{2})$ and an arrow in each Loewy diagram from the second-row factor $w(\mathcal{L}_{-1/2}^R)$ ($w(\mathcal{L}_{-3/2}^R)$) to the third-row factor $\mathcal{L}_{-3/2}^R$ ($\mathcal{L}_{-1/2}^R$).

We finish with three remarks.

- The structures of the $\mathcal{P}_{[\lambda]}^R$, for both $k = \frac{1}{2}$ and $-\frac{1}{2}$, may be partially characterised in terms of short exact sequences involving the $\mathcal{V}_{[\lambda]}^R$ of (4.31) and (4.32):

$$(4.38) \quad 0 \longrightarrow \mathcal{V}_{[\lambda+1]}^R \longrightarrow \mathcal{P}_{[\lambda]}^R \longrightarrow \mathcal{V}_{[\lambda]}^R \longrightarrow 0.$$

This unsurprising characterisation is nevertheless not as informative as the Loewy diagrams of Figure 4.

- Given experience with other “logarithmic” vertex-operator superalgebras, see [18] for example, one might be surprised by the existence of the family of logarithmic $W_{\pm 1/2}^{\min}$ -modules $\{\mathcal{P}_{[\lambda]}^R : [\lambda] \in \mathbb{C}/2\mathbb{Z}\}$. Indeed, the standard module formalism of [18, 47] suggests that the “typical” members of the family of “standard modules” should be both irreducible and projective, hence logarithmic modules should be (in this sense) rare. In the case at hand, the ubiquity of logarithmic modules is a simple consequence of the fact that $W_{\pm 1/2}^{\text{pr}}$ is not rational and, in fact, admits a logarithmic module (see [8, §4.2] for another example of this phenomenon). Examples like these indicate that the simple “typical/atypical” distinction of the standard module formalism needs generalising.
- A natural question is whether the $\mathcal{P}_{[\lambda]}^R$ are projective in a suitable category of $W_{\pm 1/2}^{\min}$ -modules. Experience with projectives for $L_k(\mathfrak{sl}_2)$ [9, 17] and (conjecturally) $L_{-3/2}(\mathfrak{sl}_3)$ [19] suggests that the answer is no because the $\mathcal{P}_{[\lambda]}^R$ are lower bounded. It would be extremely interesting to investigate the construction of logarithmic $W_{\pm 1/2}^{\min}$ -modules that do not have a lower-bounded spectral flow image. We expect that these modules will feature Jordan blocks of rank greater than 2 in the action of T_0 .




FIGURE 4. Loewy diagrams and top-space structures of the modules $\mathcal{P}_{[[\pm 1/2]]}^R$ when $k = -\frac{1}{2}$. The top row of the structures corresponds to vectors with label “ t ”, the middle row to those with labels “ m ” and “ $\bar{m}$ ”, and the bottom row to those with labels “ b ”. The shading again suggests the composition factors.

REFERENCES

- [1] D Adamović. A realization of certain modules for the $N = 4$ superconformal algebra and the affine Lie algebra $A_2^{(1)}$. *Transform. Groups*, 21:299–327, 2016. arXiv:1407.1527 [math.QA].
- [2] D Adamović. Realizations of simple affine vertex algebras and their modules: the cases $\widehat{sl}(2)$ and $\widehat{osp}(1,2)$. *Comm. Math. Phys.*, 366:1025–1067, 2019. arXiv:1711.11342 [math.QA].
- [3] D Adamović and A Babichenko. Nappi–Witten vertex operator algebra via inverse quantum hamiltonian reduction. *Comm. Contemp. Math.* (to appear), arXiv:2409.02093 [math.QA].
- [4] D Adamović, T Creutzig and N Genra. Relaxed and logarithmic modules of $\widehat{sl}_3$. *Math. Ann.*, 389:281–324, 2024. arXiv:2110.15203 [math.RT].
- [5] D Adamović, T Creutzig, N Genra and J Yang. The vertex algebras $\mathcal{R}^{(p)}$ and $\mathcal{V}^{(p)}$. *Comm. Math. Phys.*, 383:1207–1241, 2021. arXiv:2001.08048 [math.RT].
- [6] D Adamović, V Kac, P Möseneder Fradjia, P Papi and O Perše. Conformal embeddings of affine vertex algebras in minimal W-algebras I: structural results. *J. Algebra*, 500:117–152, 2018. arXiv:1602.04687 [math.RT].
- [7] D Adamović, K Kawasetsu and D Ridout. A realisation of the Bershadsky–Polyakov algebras and their relaxed modules. *Lett. Math. Phys.*, 111:38, 2021. arXiv:2007.00396 [math.QA].
- [8] D Adamović, K Kawasetsu and D Ridout. Weight module classifications for Bershadsky–Polyakov algebras. *Comm. Contemp. Math.*, 26:2340063, 2024. arXiv:2303.03713 [math.QA].
- [9] T Arakawa, T Creutzig and K Kawasetsu. Weight representations of affine Kac–Moody algebras and small quantum groups. *Adv. Math.*, 477:110365, 2025. arXiv:2311.10233 [math.RT].
- [10] T Arakawa, T Kuwabara and S Möller. Hilbert schemes of points in the plane and quasi-lisse vertex algebras with $\mathcal{N} = 4$ symmetry. arXiv:2309.17308 [math.RT].
- [11] T Arakawa and J van Ekeren. Rationality and fusion rules of exceptional W-algebras. *J. Eur. Math. Soc.*, 25:2763–2813, 2023. arXiv:1905.11473 [math.RT].

- [12] N Berkovits, C Vafa and E Witten. Conformal field theory of AdS background with Ramond–Ramond flux. *J. High Energy Phys.*, 9903:018, 1999. arXiv:hep-th/9902098.
- [13] S Berman, C Dong and S Tan. Representations of a class of lattice type vertex algebras. *J. Pure Appl. Algebra*, 176:27–47, 2002. arXiv:math.QA/0109215.
- [14] K Costello and D Gaiotto. Vertex operator algebras and 3d $\mathcal{N} = 4$ gauge theories. *J. High Energy Phys.*, 1905:18, 2019. arXiv:1804.06460 [hep-th].
- [15] T Creutzig, J Fasquel, N Genra and D Ridout. $\mathfrak{sl}_{2|1}$ minimal models I: classification of irreducible modules. In preparation.
- [16] T Creutzig, S Kanade and A Linshaw. Simple current extensions beyond semi-simplicity. *Comm. Contemp. Math.*, 22:1950001, 2020. arXiv:1511.08754 [math.QA].
- [17] T Creutzig, S Kanade, T Liu and D Ridout. Cosets, characters and fusion for admissible-level $\mathfrak{osp}(1|2)$ minimal models. *Nucl. Phys.*, B938:22–55, 2018. arXiv:1806.09146 [hep-th].
- [18] T Creutzig and D Ridout. Logarithmic conformal field theory: beyond an introduction. *J. Phys.*, A46:494006, 2013. arXiv:1303.0847 [hep-th].
- [19] T Creutzig, D Ridout and M Rupert. A Kazhdan–Lusztig correspondence for $L_{-3/2}(\mathfrak{sl}_3)$. *Comm. Math. Phys.*, 400:639–682, 2023. arXiv:2112.13167 [math.RT].
- [20] A De Sole and V Kac. Finite vs affine W -algebras. *Jpn. J. Math.*, 1:137–261, 2006. arXiv:math-ph/0511055.
- [21] M Douglas and G Moore. D-branes, quivers and ALE instantons. arXiv:hep-th/9603167.
- [22] L Eberhardt, M Gaberdiel and R Gopakumar. The worldsheet dual of the symmetric product CFT. *J. High Energy Phys.*, 1904:103, 2019. arXiv:1812.01007 [hep-th].
- [23] T Eguchi, H Ooguri and Y Tachikawa. Notes on the $K3$ surface and the Mathieu group M_{24} . *Exper. Math.*, 20:91–96, 2011. arXiv:1004.0956 [hep-th].
- [24] T Eguchi and A Taormina. Unitary representations of the $N = 4$ superconformal algebra. *Phys. Lett.*, B196:75–81, 1987.
- [25] J Fasquel, Z Fehily, E Fursman and S Nakatsuka. Connecting affine W -algebras: a case study of $\mathfrak{sl}_4$. arXiv:2408.13785 [math.QA].
- [26] J Fasquel, V Kovalchuk and S Nakatsuka. On Virasoro-type reductions and inverse hamiltonian reductions for W -algebras and W_∞ -algebras. arXiv:2411.10694 [math.QA].
- [27] J Fasquel, C Raymond and D Ridout. Modularity of admissible-level $\mathfrak{sl}_3$ minimal models with denominator 2. *Comm. Math. Phys.* (to appear), arXiv:2406.10646 [math.QA].
- [28] Z Fehily. Subregular W -algebras of type A. *Comm. Contemp. Math.*, 25:2250049, 2023. arXiv:2111.05536 [math.QA].
- [29] Z Fehily. Inverse reduction for hook-type W -algebras. *Comm. Math. Phys.*, 405:214, 2024. arXiv:2306.14673 [math.QA].
- [30] Z Fehily and D Ridout. Modularity of Bershadsky–Polyakov minimal models. *Lett. Math. Phys.*, 122:46, 2022. arXiv:2110.10336 [math.QA].
- [31] A Ferrari and A Suter. $L_1(\mathfrak{psl}_{n|n})$ from BRST reductions, associated varieties and nilpotent orbits. arXiv:2409.13028 [math.QA].
- [32] D Friedan, E Martinec and S Shenker. Conformal invariance, supersymmetry and string theory. *Nucl. Phys.*, B271:93–165, 1986.
- [33] M Gaberdiel and S Gerigk. The massless string spectrum on $\text{AdS}_3 \times S^3$ from the supergroup. *J. High Energy Phys.*, 1110:045, 2011. arXiv:1107.2660 [hep-th].
- [34] M Gaberdiel and E Mazzucchelli. The $\mathfrak{u}(2|2)_1$ WZW model. *J. Phys.*, A57:175401, 2024. arXiv:2312.03135 [hep-th].
- [35] G Götz, T Quella and V Schomerus. The WZNW model on $PSU(1, 1|2)$. *J. High Energy Phys.*, 0703:003, 2007. arXiv:hep-th/0610070.
- [36] V Kac, P Möseneder Frajria and P Papi. Spectral flow for minimal W -algebras and application to unitarity of their representations. arXiv:2508.06873 [math.RT].
- [37] V Kac, P Möseneder Frajria and P Papi. Unitarity of minimal W -algebras and their representations I. *Comm. Math. Phys.*, 401:79–145, 2023. arXiv:2208.02101 [math.RT].
- [38] V Kac, S Roan and M Wakimoto. Quantum reduction for affine superalgebras. *Comm. Math. Phys.*, 241:307–342, 2003. arXiv:math-ph/0302015.
- [39] V Kac and M Wakimoto. Quantum reduction and representation theory of superconformal algebras. *Adv. Math.*, 185:400–458, 2004. arXiv:math-ph/0304011.
- [40] V Kac and W Wang. Vertex operator superalgebras and their representations. In *Mathematical aspects of conformal and topological field theories and quantum groups*, volume 175 of *Contemporary Mathematics*, pages 161–191, Providence, 1994. American Mathematical Society. arXiv:hep-th/9312065.
- [41] H Kausch. Symplectic fermions. *Nucl. Phys.*, B583:513–541, 2000. arXiv:hep-th/0003029.
- [42] K Kawasetsu and D Ridout. Relaxed highest-weight modules I: rank 1 cases. *Comm. Math. Phys.*, 368:627–663, 2019. arXiv:1803.01989 [math.RT].
- [43] H Li. The physics superselection principle in vertex operator algebra theory. *J. Algebra*, 196:436–457, 1997.
- [44] H Nakajima. Instantons on ALE spaces, quiver varieties, and Kac–Moody algebras. *Duke Math. J.*, 76:365–416, 1994.
- [45] H Nakajima. Quiver varieties and kac-moody algebras. *Duke Math. J.*, 91:515–560, 1998.
- [46] D Ridout. $\widehat{\mathfrak{sl}}(2)_{-1/2}$: A case study. *Nucl. Phys.*, B814:485–521, 2009. arXiv:0810.3532 [hep-th].
- [47] D Ridout and S Wood. The Verlinde formula in logarithmic CFT. *J. Phys. Conf. Ser.*, 597:012065, 2015. arXiv:1409.0670 [hep-th].
- [48] A Semikhatov. Inverting the Hamiltonian reduction in string theory. In *28th International Symposium on Particle Theory, Wendisch-Rietz, Germany*, pages 156–167, 1994. arXiv:hep-th/9410109.
- [49] K Thielemans. A MATHEMATICA package for computing operator product expansions. *Int. J. Mod. Phys.*, C2:787–798, 1991.
- [50] J Troost. Massless particles on supergroups and $\text{AdS}_3 \times S^3$ supergravity. *J. High Energy Phys.*, 1107:042, 2011. arXiv:1102.0153 [hep-th].

[51] Y Zhu. Modular invariance of characters of vertex operator algebras. *J. Amer. Math. Soc.*, 9:237–302, 1996.

(Zachary Fehily) SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, PARKVILLE, AUSTRALIA, 3010.

Email address: zacfehily@gmail.com

(Christopher Raymond) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAMBURG, HAMBURG, 20146, GERMANY.

Email address: christopher.raymond@uni-hamburg.de

(David Ridout) SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, PARKVILLE, AUSTRALIA, 3010.

Email address: david.ridout@unimelb.edu.au