

347 FINAL, IN CLASS PART
DECEMBER 12, 2005

(1) Let $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ be (well-defined) maps. Complete the following sentences:

(a) (4 points) By definition, f is surjective if and only if

(b) (4 points) f is not surjective if and only if

(c) (11 points) Prove: If g is not surjective, then neither is $g \circ f$.

- (2) Let X be a set, and let P and Q be properties that elements of X could have. We write $P(x)$ for “ x has the property P ”, and $Q(x)$ for “ x has the property Q ”. Consider the sets

$$Y := \{y \in X \mid P(y)\}$$

and

$$Z := \{x \in X \mid Q(x)\}.$$

Complete the following sentences **with statements about the properties P and Q** :

(a) (3 points) $Y = Z$ if and only if

(b) (3 points) $x \in X \setminus Y$ if and only if

(c) (3 points)

$$Y \setminus Z = \{x \in X \mid \quad \quad \quad \}$$

(d) (3 points) $Y = X \setminus Z$ if and only if

(e) (3 points) $Y \cup Z = Y$ if and only if

- (3) (2 points for every correct answer and -1 point for every false answer) For each of the following statements, indicate whether it is equivalent to the negation of $A \Rightarrow B$.
- (a) **Yes No** A is true and B is not true
 - (b) **Yes No** (not A) or B
 - (c) **Yes No** if A holds then not B holds.
 - (d) **Yes No** A is necessary for B .
 - (e) **Yes No** If B holds then A does not hold.

- (4) **Permutations:** Consider the permutation f with two-line form

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 2 & 7 & 3 & 6 & 4 & 5 \end{pmatrix}$$

- (a) (3 points) Write f in cycle form.
- (b) (2 points) What is the order of f ?
- (c) (3 points) Compute f^{-1} in cycle or in two-line form.

(d) (5 points) Compute $f^{-1} \circ f^{-1}$ in cycle or in two-line form.

(e) (3 points) What is $f \circ f \circ f \circ f \circ f \circ f \circ f \circ f \circ f \circ f$?

(5) **Modular arithmetic:**

(a) (4 points) Compute $7^{29000002}$ modulo 59.

(b) (8 points) What is the multiplicative inverse of $[12]$ in $\mathbb{Z}/139\mathbb{Z}$?

- (6) (11 points) How would you structure a proof of the following statement about a function $f: \mathbb{R} \rightarrow \mathbb{R}$?

For every $\varepsilon > 0$, there exists a $\delta > 0$ such that for every $x, y \in \mathbb{R}$ the following holds:

$$|x - y| < \delta \Rightarrow |f(x) - f(y)| < \varepsilon.$$

Remark: At some point in this proof you would have to make a clever guess for something. Just write “???” in the place of what the guess would have to be.

- (7) (6 points for a correct answer, -2 points for a false answer) Compare the statement of the previous question to the one from Midterm 3:

For every $x_0 \in \mathbb{R}$ and every $\varepsilon > 0$, there exists a $\delta > 0$ such that for every $x \in \mathbb{R}$ the following holds:

$$|x - x_0| < \delta \Rightarrow |f(x) - f(x_0)| < \varepsilon.$$

Which of the two do you think is the stronger statement?
Explain why.

(8) (11 points) Use induction, starting with $n = 0$, to prove: For every $n \in \mathbb{N}$,

$$\sum_{j=1}^n j = \frac{n(n+1)}{2}.$$